

PCT
INTERNATIONAL COOPERATION TREATY

PCT

NOTIFICATION OF THE RECORDING
OF A CHANGE(PCT Rule 92bis.1 and
Administrative Instructions, Section 422)

From the INTERNATIONAL BUREAU

To:

WALKER, Naomi
Kiryat Weizmann Science Park
P.O. Box 2313
76123 Rehovot
ISRAËL

Date of mailing (day/month/year) 27 November 2000 (27.11.00)	IMPORTANT NOTIFICATION
Applicant's or agent's file reference DPL/009/PCT	
International application No. PCT/IL00/00562	International filing date (day/month/year) 13 September 2000 (13.09.00)

1. The following indications appeared on record concerning:

☒ the applicant

 ☒ the inventor

 ☐ the agent

 ☐ the common representative

Name and Address	State of Nationality	State of Residence
	Telephone No.	
	Facsimile No.	
	Teleprinter No.	

2. The International Bureau hereby notifies the applicant that the following change has been recorded concerning:

☒ the person

 ☒ the name

 ☒ the address

 ☒ the nationality

 ☒ the residence

Name and Address AYALON, Oran Shikun Banim 143 Ganei Yohanan 76922 Israel	State of Nationality IL	State of Residence IL
	Telephone No.	
	Facsimile No.	
	Teleprinter No.	

3. Further observations, if necessary:

Applicant/inventor has been added.

4. A copy of this notification has been sent to:

<input checked="" type="checkbox"/> the receiving Office	<input type="checkbox"/> the designated Offices concerned
<input checked="" type="checkbox"/> the International Searching Authority	<input type="checkbox"/> the elected Offices concerned
<input type="checkbox"/> the International-Preliminary Examining Authority	<input type="checkbox"/> other:

The International Bureau of WIPO 34, chemin des Colombettes 1211 Geneva 20, Switzerland Facsimile No.: (41-22) 740.14.35	Authorized officer Athina Nickitas-Etienne Telephone No.: (41-22) 338.83.38
---	---

PATENT COOPERATION TREATY

PCT

NOTIFICATION CONCERNING
AMENDMENTS OF THE CLAIMS

(PCT Rule 62 and
Administrative Instructions, Section 417)

From the INTERNATIONAL BUREAU

To:

Commissioner
US Department of Commerce
United States Patent and Trademark
Office, PCT
2011 South Clark Place Room
CP2/5C24
Arlington, VA 22202
ETATS-UNIS D'AMERIQUE
in its capacity as International Preliminary Examining Authority

Date of mailing (day/month/year)

22 October 2001 (22.10.01)

International application No.

PCT/IL00/00562

International filing date (day/month/year)

13 September 2000 (13.09.00)

Applicant

D-PHARM LIMITED et al

The International Bureau hereby informs the International Preliminary Examining Authority that no amendments under Article 19 have been received by the International Bureau (Administrative Instructions, Section 417).

The International Bureau of WIPO
34, chemin des Colombettes
1211 Geneva 20, Switzerland

Facsimile No. (41-22) 740.14.35

Authorized officer

Ali SOLEIMAN

Telephone No. (41-22) 338.83.38

PCT COOPERATION TREATY

PCT

NOTIFICATION OF RECEIPT OF RECORD COPY

(PCT Rule 24.2(a))

From the INTERNATIONAL BUREAU

To:

WALKER, Naomi
Kiryat Weizmann Science Park
P.O. Box 2313
76123 Rehovot
ISRAËL

Date of mailing (day/month/year) 26 October 2000 (26.10.00)	IMPORTANT NOTIFICATION
Applicant's or agent's file reference DPL/009/PCT	International application No. PCT/IL00/00562

The applicant is hereby notified that the International Bureau has received the record copy of the international application as detailed below.

Name(s) of the applicant(s) and State(s) for which they are applicants:

D-PHARM LIMITED (for all designated States except US)
KOZAK, Alexander et al (for US)

International filing date : 13 September 2000 (13.09.00)
Priority date(s) claimed : 14 September 1999 (14.09.99)
Date of receipt of the record copy
by the International Bureau : 03 October 2000 (03.10.00)
List of designated Offices :

AP : GH,GM,KE,LS,MW,MZ,SD,SL,SZ,TZ,UG,ZW
EA : AM,AZ,BY,KG,KZ,MD,RU,TJ,TM
EP : AT,BE,CH,CY,DE,DK,ES,FI,FR,GB,GR,IE,IT,LU,MC,NL,PT,SE
OA : BF,BJ,CF,CG,CI,CM,GA,GN,GW,ML,MR,NE,SN,TD,TG
National : AE,AG,AL,AM,AT,AU,AZ,BA,BB,BG,BR,BY,BZ,CA,CH,CN,CR,CU,CZ,DE,DK,DM,DZ,EE,
ES,FI,GB,GD,GE,GH,GM,HR,HU,ID,IL,IN,IS,JP,KE,KG,KP,KR,KZ,LC,LK,LR,LS,LT,LU,LV,MA,
MD,MG,MK,MN,MW,MX,MZ,NO,NZ,PL,PT,RO,RU,SD,SE,SG,SI,SK,SL,TJ,TM,TR,TT,TZ,UA,UG,US,
UZ,VN,YU,ZA,ZW

ATTENTION

The applicant should carefully check the data appearing in this Notification. In case of any discrepancy between these data and the indications in the international application, the applicant should immediately inform the International Bureau.

In addition, the applicant's attention is drawn to the information contained in the Annex, relating to:

- ☒ time limits for entry into the national phase
☐ confirmation of precautionary designations
☐ requirements regarding priority documents

A copy of this Notification is being sent to the receiving Office and to the International Searching Authority.

The International Bureau of WIPO 34, chemin des Colombettes 1211 Geneva 20, Switzerland Facsimile No. (41-22) 740.14.35	Authorized officer: Athina Nickitas-Etienne Telephone No. (41-22) 338.83.38
--	---

PCT
ENT COOPERATION TREA

PCT

NOTIFICATION OF THE RECORDING
OF A CHANGE(PCT Rule 92bis.1 and
Administrative Instructions, Section 422)

From the INTERNATIONAL BUREAU

To:

WALKER, Naomi
Kiryat Weizmann Science Park
P.O. Box 2313
76123 Rehovot
ISRAËLDate of mailing (day/month/year)
27 November 2000 (27.11.00)Applicant's or agent's file reference
DPL/009/PCT

IMPORTANT NOTIFICATION

International application No.
PCT/IL00/00562International filing date (day/month/year)
13 September 2000 (13.09.00)

1. The following indications appeared on record concerning:

☒ the applicant ☒ the inventor ☐ the agent ☐ the common representative

Name and Address

SENDERICHIN, Alexander
Eben Ezra 11/32
77622 Ashdod
Israel

State of Nationality

IL

State of Residence

IL

Telephone No.

Facsimile No.

Teleprinter No.

2. The International Bureau hereby notifies the applicant that the following change has been recorded concerning:

☐ the person ☒ the name ☐ the address ☐ the nationality ☐ the residence

Name and Address

SENDERIKHIN, Alexander
Eben Ezra 11/32
77622 Ashdod
Israel

State of Nationality

IL

State of Residence

IL

Telephone No.

Facsimile No.

Teleprinter No.

3. Further observations, if necessary:

4. A copy of this notification has been sent to:

☒ the receiving Office ☐ the designated Offices concerned
☒ the International Searching Authority ☐ the elected Offices concerned
☐ the International Preliminary Examining Authority ☐ other:The International Bureau of WIPO
34, chemin des Colombettes
1211 Geneva 20, Switzerland

Facsimile No.: (41-22) 740.14.35

Authorized officer

Athina Nickitas-Etienne

Telephone No.: (41-22) 338.83.38

PCT
INTERNATIONAL COOPERATION TREATY

PCT

NOTIFICATION OF THE RECORDING
OF A CHANGE(PCT Rule 92bis.1 and
Administrative Instructions, Section 422)

From the INTERNATIONAL BUREAU

To:

WALKER, Naomi
Kiryat Weizmann Science Park
P.O. Box 2313
76123 Rehovot
ISRAËL

Date of mailing (day/month/year) 27 November 2000 (27.11.00)	IMPORTANT NOTIFICATION
Applicant's or agent's file reference DPL/009/PCT	
International application No. PCT/IL00/00562	International filing date (day/month/year) 13 September 2000 (13.09.00)

1. The following indications appeared on record concerning:

☒ the applicant

 ☐ the inventor

 ☐ the agent

 ☐ the common representative

Name and Address D-PHARM LIMITED Kiryat Weizmann Science Park P.O. Box 2313 76326 Rehovot Israel	State of Nationality IL	State of Residence IL
	Telephone No.	
	Facsimile No.	
	Teleprinter No.	

2. The International Bureau hereby notifies the applicant that the following change has been recorded concerning:

☐ the person

 ☐ the name

 ☒ the address

 ☐ the nationality

 ☐ the residence

Name and Address D-PHARM LIMITED Kiryat Weizmann Science Park P.O. Box 2313 76123 Rehovot Israel	State of Nationality IL	State of Residence IL
	Telephone No.	
	Facsimile No.	
	Teleprinter No.	

3. Further observations, if necessary:

4. A copy of this notification has been sent to:

<input checked="" type="checkbox"/> the receiving Office	<input type="checkbox"/> the designated Offices concerned
<input checked="" type="checkbox"/> the International Searching Authority	<input type="checkbox"/> the elected Offices concerned
<input type="checkbox"/> the International Preliminary Examining Authority	<input type="checkbox"/> other:

The International Bureau of WIPO 34, chemin des Colombettes 1211 Geneva 20, Switzerland Facsimile No.: (41-22) 740.14.35	Authorized officer Athina Nickitas-Etienne Telephone No.: (41-22) 338.83.38
--	--

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
22 March 2001 (22.03.2001)

PCT

(10) International Publication Number
WO 01/19320 A2

- (51) International Patent Classification⁷: **A61K**
- (21) International Application Number: **PCT/IL00/00562**
- (22) International Filing Date:
13 September 2000 (13.09.2000)
- (25) Filing Language: **English**
- (26) Publication Language: **English**
- (30) Priority Data:
131887 14 September 1999 (14.09.1999) **IL**
- (71) Applicant (for all designated States except US):
D-PHARM LIMITED [IL/IL]; Kiryat Weizmann
Science Park, P.O. Box 2313, 76123 Rehovot (IL).

- Eben Ezra 11/32, 77622 Ashdod (IL). **AYALON, Oran**
[IL/IL]; Shikun Banim 143, 76922 Ganei Yohanan (IL).
- (74) Agent: **WALKER, Naomi**; Kiryat Weizmann Science
Park, P.O. Box 2313, 76123 Rehovot (IL).
- (81) Designated States (*national*): AE, AG, AL, AM, AT, AU,
AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CR, CU, CZ,
DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR,
HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR,
LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ,
NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM,
TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.
- (84) Designated States (*regional*): ARIPO patent (GH, GM,
KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian
patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European
patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE,
IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG,
CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

- (72) Inventors; and
- (75) Inventors/Applicants (for US only): **KOZAK, Alexan-**
der [IL/IL]; 8 Naphtali-Ben-Ephraim Street, 76555 Re-
hovot (IL). **SHAPIRO, Israel** [IL/IL]; 1 Aharon Bouga-
nium Street, 75122 Ramla (IL). **VINNIKOVA, Marina**
[IL/IL]; 1 Aharon Bouganium Street, 75122 Ramla (IL).
ERSHOV, Leonid [IL/IL]; Ganei Shemesh 64/20, 99510
Beit Shemesh (IL). **SENDERIKHIN, Alexander** [IL/IL];

Published:

— Without international search report and to be republished
upon receipt of that report.

For two-letter codes and other abbreviations, refer to the "Guid-
ance Notes on Codes and Abbreviations" appearing at the begin-
ning of each regular issue of the PCT Gazette.

(54) Title: **PHOSPHOLIPID PRODRUGS OF ANTI-PROLIFERATIVE DRUGS**

(57) Abstract: The invention discloses prodrugs comprising anti-proliferative drugs covalently linked, via a bridging group, to a phospholipid moiety such that the active species is preferentially released, preferably by enzymatic cleavage, at the required site of action. The invention further discloses pharmaceutical compositions comprising said prodrugs and the uses thereof for the treatment of diseases and disorders related to inflammatory, to degenerative or atrophic conditions, and to uncontrolled cell growth.



WO 01/19320 A2

PHOSPHOLIPID PRODRUGS OF ANTI-PROLIFERATIVE DRUGS

5 FIELD OF THE INVENTION

The present invention relates to prodrugs of anti-proliferative agents comprising a phospholipid moiety covalently linked, via a bridging group, to an anti-proliferative agent, such that the active species is preferentially released, preferably by enzymatic cleavage, at the required site of action.

10 The invention further relates to pharmaceutical compositions comprising said prodrugs and to the use thereof for the treatment of diseases and disorders related to inflammatory, degenerative or atrophic conditions and to uncontrolled cell growth.

BACKGROUND OF THE INVENTION

15 Anti-proliferative drugs, also known as anti-metabolites, act by inhibiting crucial metabolic processes and are commonly used in the treatment of diseases involving abnormal cell proliferation, such as tumors. Many anti-proliferative drugs are also useful as anti-inflammatory agents, exerting their effect by suppressing local or even systemic inflammatory responses mediated by the immune system. However,
20 the utility of these drugs is severely hampered by their excessive toxicity and adverse side effects on healthy cells of the treated patient.

Methotrexate (MTX) is an effective anti-proliferative drug commonly used in cancer therapy. MTX is also known as a leading anti-inflammatory drug, and is currently the main drug used in the treatment of rheumatoid arthritis (RA). It is the
25 drug of first choice for treating children with recalcitrant juvenile RA.

Methotrexate is an analogue of dihydrofolate that inhibits the enzyme dihydrofolate reductase (DHFR), thus depleting intracellular tetrahydrofolate (FH₄) which is an essential co-factor required for the *de novo* synthesis of purine nucleotides.

30 Another anti-metabolite widely used in cancer chemotherapy is the pyrimidine analog fluorodeoxyuridine (FUdR). FUdR is converted *in vivo* into

fluorodeoxyuridylate (F-dUMP) which is an analog of the normal substrate dUMP and irreversibly inhibits thymidylate synthase.

MTX and FUdR are valuable drugs in the treatment of many rapidly growing tumors, however, their use is limited by the frequency and severity of side effects.

- 5 Unwanted side effects include toxicity to all rapidly dividing normal cells, such as stem cells in the bone marrow, epithelial cells of the intestinal tract, etc. Folic acid antagonists are also toxic to developing embryos. Treatment with MTX is especially problematic in patients having chronic debilitating inflammatory diseases that require prolonged therapy, such diseases as rheumatoid arthritis, asthma, dermatological
- 10 diseases such as psoriasis and gastrointestinal inflammation such as Crohn's disease. These patients may suffer from induced nephrotoxicity, due to precipitation of the drug or its metabolites in the renal tubes, and from hepatic fibrosis and cirrhosis.

- Another major problem in chemotherapy, which is particularly relevant in the case of anti-metabolites, is inherent or acquired resistance of tumors to cytotoxic
- 15 drugs. For example, development of resistance to MTX frequently follows prolonged exposure to this drug. Resistance may be due to new mutations induced by the clinical treatment or to positive selection, by the treatment regimen, of pre-existing resistant mutant cell. Known mechanisms for development of resistance involve impaired transport of MTX into cells, e.g. by mutations in the Reduced Folates
- 20 Carrier (RFC), over expression of the target enzyme DHFR, or mutations in the enzyme responsible for polyglutamination of reduced folates (FPGS).

- A more severe problem in the clinic is the phenomenon of multi-drug resistance (MDR), which is a resistance to a broad spectrum of structurally unrelated cytotoxic drugs. MDR is mediated by membranal "pumps" which actively expel
- 25 chemotherapeutic drugs from the tumor cells. Two drug pumps commonly found in cancer are P-glycoprotein (P-gp) and multidrug resistance-associated protein (MRP). MDR significantly limits the efficacy of many cancer chemotherapy regimens and is a major factor in the failure of cancer chemotherapy.

- It would, therefore, be most advantageous to have drug derivatives that are
- 30 targeted or selectively active in the diseased cells rather than in the healthy cells, thus reducing undesirable side effects. It would also be desirable to generate new anti-

proliferative agents that overcome drug-resistance, as well as agents that are active as cytotoxic drugs but lack or have a reduced ability to provoke MDR phenotype.

The use of prodrugs to impart desired characteristics such as increased bioavailability or increased site-specificity for known drugs is a recognized concept in the state of the art of pharmaceutical development. The use of various lipids in the preparation of particular types of prodrugs is also known in the background art. In none of those instances are the prodrugs able to achieve preferential accumulation of an anti-proliferative drug within the diseased cells of the organ, by activation with intracellular lipases. Rather, the prodrugs enable the drug to be transported to a specific site, or to be released within a specific organ, as in the case of the phospholipid prodrugs of salicylates and non-steroidal anti-inflammatory drugs disclosed in International Patent Application WO 91/16920 which, taken orally, protect the gastric mucosa and release the active principle in the gut.

In other examples of phospholipid prodrugs, the formulation of the prodrugs into liposomes or other micellar structures is the feature that enables their preferential uptake, for instance by liver cells or by macrophages as in the case of the phospholipid conjugates of antiviral drugs disclosed in International patent applications WO 93/00910 and WO 90/00555.

U.S Patent No. 5,411,947 discloses a method of converting a drug to an orally available form by covalently bonding a lipid to the drug. It discloses a lipid prodrug of 5-fluorouridine, but differs from the present invention in that the drug is attached to the lipid at the R³ position.

U.S Patent No. 4,772,594 discloses prodrugs containing 5-Fluorouracil, but differs from the present invention in that the drug is attached to cholesterol.

U.S. Patent Nos. 5,149,794 and 5,543,389 disclose covalent polar lipid-drug conjugates to facilitate the entry of drugs into cells at pharmacokinetically useful levels. In contrast to the present invention, the disclosed prodrugs are directed to certain intracellular structures and organelles due to the existence of the polar lipid carrier, and drug release from the lipid conjugate is not a requirement for the drug targeting. Moreover, the lipid-drug conjugates may or may not include a spacer and it is explicitly stated that the conjugates may be pharmacologically active themselves.

Though phospholipids are included in a list of potentially useful polar lipids, the drug-phospholipid conjugates were clearly never reduced to practice and have not been suggested to be inactive derivatives of the drug.

5 Mono- and di-esters of methotrexate having short alkyl chains ranging from one to four carbons in length, and MTX mono-esters having alkyl chains of up to sixteen carbon atoms, and their cytotoxic effects on cultured cells *in vitro* have been disclosed by Rosowsky et. (J. Med. Chem., 1978, 21: 380-386; J. Med. Chem., 1984, 27: 605-609). However, none of the disclosed compounds is a phospholipid derivative of MTX as in the present invention.

10 Non-lipid analogues of methotrexate have been previously disclosed (for example by Antonjuk, D. et al., 1984 J. Chem. Soc. Perkin Trans. 1 (9) 1989-2003), however the derivatives are monoamides and not phospholipids as in the present invention.


15 International Application WO 94/22483 describes prodrugs which selectively release pharmacologically active compounds in hyperactivated cells, but does not disclose methotrexate or 5-fluorodeoxyuridine derivatives.

SUMMARY OF THE INVENTION

20 The object of the present invention is to provide agents with improved anti-proliferative properties. It is a further object of the present invention to provide prodrugs of anti-proliferative drugs that undergo preferential activation within the disease-affected cells and tissues.

25 The prodrugs in accordance with the present invention comprise a phospholipid moiety covalently linked, via a bridging group, to an anti-proliferative drug residue, such that the active species is preferentially released, preferably by enzymatic cleavage, at the required site of action.

Thus, the present invention provides, in a first aspect, a prodrug of the general formula I:



Formula I

or a pharmaceutically acceptable salt thereof, wherein:

- 5 **R1** is a saturated or unsaturated, straight-chain or branched, substituted or unsubstituted hydrocarbon chain having from 2 to 30 carbon atoms;
R2 is H or a phospholipid head group;
Z is saturated or unsaturated, straight-chain or branched, substituted or unsubstituted hydrocarbon chain having from 2 to 15 carbon atoms, which may include cyclic
10 elements, and optionally is interrupted by one or more atoms selected from oxygen and sulfur atoms;
X is a direct covalent bond or selected from the group consisting of O, S, NH and C(O) groups; and
D is a residue of an anti-proliferative drug,
15 wherein the bound anti-proliferative drug residue is an inactive form of the drug which is selectively activated in cells and tissues with elevated phospholipase activity.

20 According to a currently preferred embodiment, R1 of the prodrug of the general formula I is a hydrocarbon chain having from 5 to 20 carbon atoms, more preferably 15 or 17 carbon atoms.

In another preferred embodiment R2 of the prodrug of the general formula I is selected from the group consisting of choline, ethanolamine, inositol and serine. Preferred anti-proliferative drugs used in the prodrug of the general formula I are methotrexate and 2'-deoxy-5-fluorouridine.

Currently more preferred prodrugs according to the invention are selected from the group consisting of:

1-Stearoyl-2-[3-(α -MTX amido)-Propanoyl]-sn-Glycero-3-Phosphatidylcholine,

5 1-Stearoyl-2-[3-(γ -dodecylate- α -MTX amido)-Propanoyl]-sn-Glycero-3-Phosphatidylcholine,

1-Stearoyl-2-[4-(α -MTX amido)-Butanoyl]-sn-Glycero-3-Phosphatidylcholine,

10 1-Stearoyl-2-[6-(α -MTX amido)-Hexanoyl]-sn-Glycero-3-Phosphatidylcholine,

1-Stearoyl-2-[8-(α -MTX amido)-Octanoyl]-sn-Glycero-3-Phosphatidylcholine,

1-Stearoyl-2-[8-(γ -dodecylate- α -MTX amido)-Octanoyl]-sn-Glycero-3-Phosphatidylcholine,

15 1-Stearoyl-2-[3-(α -dodecylate- γ -MTX amido)-Propanoyl]-sn-Glycero-3-Phosphatidylcholine, and

1-stearoyl-2-[5^{''}-(2^{''}-deoxy-5'-fluorouridine-5^{''})-3^{'''},3^{'''}-dimethyl] glutaroyl-1^{'''}-sn-glycero-3-phosphatidylcholine.

Currently most preferred prodrugs are:

20 1-Stearoyl-2-[3-(α -MTX amido)-Propanoyl]-sn-Glycero-3-Phosphatidylcholine, and
1-Stearoyl-2-[3-(α -dodecylate- γ -MTX amido)-Propanoyl]-sn-Glycero-3-Phosphatidylcholine.

25 In another aspect of the invention, there is provided a pharmaceutical composition comprising a pharmaceutically acceptable carrier and, as an active ingredient, a prodrug of the general formula I as defined above.

The pharmaceutical composition, in accordance with the invention, may further include one or more additional anti-neoplastic agents.

In a further aspect, the present invention provides the use of a prodrug of the aforementioned formula I, or a pharmaceutically acceptable salt thereof, for the preparation of a pharmaceutical composition.

In still a further aspect, the invention provides methods for treatment of
5 diseases or disorders related to an inflammatory condition such as granulomatous diseases, arthritis, rheumatoid arthritis, multiple sclerosis, systemic sclerosis, asthma, psoriasis, systemic lupus erythematosus, inflammatory bowel syndromes and migraines. Also provided are methods for treatment of diseases and disorders related to a degenerative or atrophic condition, in particular central or peripheral neurological
10 diseases or disorders. The diseases and disorders related to a degenerative or atrophic condition may be selected from, but are not limited to, the group consisting of autoimmune, cerebrovascular and neurodegenerative diseases and disorders such as idiopathic dementia, vascular dementia, multi-infarct dementia, traumatic dementia, Alzheimer's disease, Pick's disease, Huntington's disease, dementia paralitica,
15 Parkinson's disease, diabetic neuropathy, amyotrophic lateral sclerosis, ischemia of the optic nerve, age-related macular degeneration, stroke and trauma. Also provided are methods for treatment of diseases or disorders related to uncontrolled cell growth. Said diseases or disorders may be selected from, but are not limited to, psoriasis, lymphocytic leukemia, myelogenous leukemia, Burkitt's
20 lymphoma, non-Hodgkin's lymphomas, mycosis fungoides, osteosarcoma, hydatidiform mole, trophoblastic diseases such as chorioadenoma destruens and choriocarcinoma, and carcinomas of the head and neck, breast, liver, lung, colon, ovary, cervix, urinary, bladder, prostate, pancreas, skin, the gastrointestinal tract and the oropharyngeal areas.

25 The aforementioned methods comprise administering to a patient in need thereof a pharmaceutical composition containing a therapeutically effective amount of a prodrug of the general formula I, in accordance with the invention.

In a preferred embodiment, the prodrugs of the invention are useful for the treatment of neoplastic growths. The neoplastic growths may be primary or secondary
30 tumors, including drug-resistant tumor, for example tumors resistant to methotrexate and multidrug-resistant tumors.

BRIEF DESCRIPTION OF THE DRAWING

Fig. 1 depicts a graph of animal survival during the course of an experiment wherein mice were i.p. transplanted with L1210 mouse leukemia cells and then treated with vehicle only (squares), MTX (triangles) or molar equivalent dose of DP-MTX-71 (circles) according to the regimen described in Example 11.

DETAILED DESCRIPTION OF THE INVENTION

The prodrugs of the general formula I according to the present invention comprise anti-proliferative agents covalently conjugated, via a bridging group, to position sn-2 of a phospholipid. These compounds, being hydrophobic in nature, may penetrate biological membranes, thus effectively transporting the prodrug into cells or organs.

According to the principles of the invention, the bound anti-proliferative drug residue is an inactive form of the drug and is linked to the phospholipid moiety through a bridging group that is susceptible to cleavage. The cleavage enables release of an active anti-proliferative drug from the prodrug compound, preferably at the required site of action. The specificity of the drug activation is afforded by a linking group that is designed to be sensitive to cleavage by lipases, preferably phospholipases (e.g. PLA₂) that are specifically elevated at the pathological target tissue. According to this process, denoted as Regulated Activation of Prodrugs (RAP), regulated release of the active drug preferentially occurs at the disease site. It should be appreciated that the release of the active anti-proliferative drug from the prodrug may be enzymatically or non-enzymatically executed. Moreover, the released active drug may be different from the original parent drug molecule, including drug derivatives wherein one or more chemical group has been removed from or added to the original anti-proliferative residue D. These cases are also included within the concept of the invention provided that the released drug derivative preserves therapeutic activity. Preferably the cleavage process of the molecule of the invention is initiated specifically at the diseased cells, thus generating a highly specific and highly effective drug release at the desired target site. However, it may not be excluded that under certain pathological conditions, the prodrug molecule may be

therapeutically active while the anti-proliferative agent is not cleaved from the phospholipid moiety.

Irrespective of the exact composition of the active component and without wishing to be limited to a particular mechanism of action, it is evident that the novel
5 prodrugs of the invention have an improved therapeutic profile. They also may enable extending the usefulness of anti-proliferative agents to conditions that do not respond to equivalent doses of the parent drug, or permit use of the drugs in certain conditions and diseases where the toxicity of the parent molecule is prohibitive.

While screening the compounds of the invention for their cytotoxic effect on
10 various tumor cell lines, it was unexpectedly found that a currently most preferred compound according to the principles of the invention, denoted herein as DP-MTX-71 showed a remarkable anti-proliferative effect on drug-resistant cells. Thus it was established that certain compounds of the invention may be useful as anti-cancer agents, inhibiting the growth of sensitive as well as resistant tumor cell lines.

15 In accordance with the principles of the present invention, the various prodrug molecules may be specifically tailored to suit specific target sites and specific indications. In designing a prodrug compound in accordance with the invention, the specific nature of the pathological condition to be treated should be considered. This involves determining the desired pharmacological activity to be achieved, hence the
20 choice of the anti-proliferative drug residue D. In addition, the phospholipid moiety may be modified in order to optimize lipophilicity of the prodrug and drug release.

The desired lipophilicity of the molecule depends on the particular site where the pharmaceutical activity is needed. Accordingly, the number of carbon atoms in the R1 hydrocarbon chain of a prodrug of the general formula I, is determined. The
25 lipophilicity of the molecule is directly correlated to the selected hydrocarbon chain length. R1 chains according to the invention may contain 2 to 30 carbon atoms. Molecules with R1 having from 5 to 20 carbon atoms are most desirable as endowing the molecule with suitable hydrophobicity for crossing biological membranes and at the same time providing adequate substrate for the action of phospholipase.
30 R1 may be a straight-chained or branched, saturated or unsaturated hydrocarbon chain, containing one or more double and/or triple bonds. One or more hydrogen atoms on

the chain may be substituted, for example, by halogen atoms or by a small alkyl group such as methyl residues, with the proviso that the substituents still allow free access for the desired cleaving enzymes. In preferred embodiments of the invention R1 is an alkyl residue of an odd number of carbon atoms. More preferably R1 is an alkyl residue of 15 or 17 carbon atoms yielding, respectively, the naturally occurring palmitoyl (C₁₆) or stearoyl (C₁₈) residues at the sn-1 position of the phospholipid.

The lipophilicity of the lipid molecule is also affected by the nature of the phospholipid head group, denoted as R2 in the prodrug of the general formula I. The phospholipid moiety may be selected from, but is not limited to, the group consisting of phosphatidic acid, phosphatidylcholine, phosphatidylethanolamine, phosphatidylinositol and phosphatidylserine.

The anti-proliferative drug residue incorporated into the phospholipid prodrug may be any anti-proliferative agent that inhibits, directly or indirectly, cell growth. The drug is covalently linked, via a bridging group, to position sn-2 of the phospholipid, such that the anti-proliferative activity of the bound drug residue is significantly reduced.

In one particular embodiment, prodrugs of the invention are phospholipid derivatives of methotrexate. In a preferred embodiment, binding of the drug to the lipid-bridge moiety is specifically directed to the α -carboxylic group of methotrexate (herein denoted α -MTX). It was found that binding of phospholipid to methotrexate via its alpha carboxylic group yields a pharmaceutically inactive methotrexate. In another preferred embodiment, the methotrexate moiety is linked to the phospholipid through the γ -carboxyl of methotrexate (herein denoted γ -MTX).

The hydrogen atom of the carboxylic group of methotrexate that is not attached to the phospholipid, i.e. the γ -carboxylic group of α -MTX or the α -carboxylic group of γ -MTX, may be substituted by a saturated or unsaturated, substituted or unsubstituted hydrocarbon chain having from 2 to 30 carbon atoms.

Preferably, the appropriate drugs are linked to the bridging group through a carboxyl, oxy, amine or mercapto group, thus generating an ester, amido or a thio bond.

Suitable anti-proliferative drugs may include, but are not limited to, anti-metabolites such as paclitaxel, estramustine, melphalan, carmustine, nimustine, daunorubicin, doxorubicin, denopterin, methotrexate, trimetrexate, fluorouracil, fluorodeoxyuridine, 6-azauridine, flutamide, nilutamide, goserelin, leuprolide and anthralin.

In currently preferred embodiments of the invention the anti-proliferative drug is selected from methotrexate and 2'-deoxy-5-fluorouridine and pharmaceutically acceptable derivatives thereof.

The anti-proliferative drug residue, D, is covalently linked to the phospholipid moiety via the bridging group $-C(O)-Z-X$. The choice of the preferred bridging group is dependent on several considerations; it should participate in a stable covalent bond with the D moiety while lending itself to cleavage at the target site. A preferred bridging group, is such that is resistant to cleavage under normal physiological conditions encountered by the administered compound on its way to the target site.

The bridging group should not confer a steric hindrance on the enzymatic cleavage of the ester bond at position sn-2 of the phospholipid of the general formula I.

Depending on the treated pathological condition and the particular diseased cell or organ, it will be desirable at times to choose such a bridging group that will regulate the release of the active drug by facilitating or delaying its cleavage from the prodrug molecule.

Component Z of the bridging group may be a saturated or unsaturated, straight-chain or branched hydrocarbon chain having from 2 to 15 carbon atoms, which may include cyclic elements, and optionally is interrupted by one or more atoms selected from oxygen and sulfur atoms. Component X may be a covalent bond or is selected from the group consisting of O, S, NH and C(O) groups. According to a preferred embodiment, the total number of carbon atoms in the bridging group $C(O)-Z-X$ is at least 4 but at most 15. It was found that this length of carbon chain provides a spacer which enables good access to an enzyme, preferably phospholipase and in particular PLA_2 , for digesting the ester bond at position sn-2 of the phospholipid of the general formula I. Shorter spacers, in particular bridging groups comprising less than three carbon atoms, may be problematic, by creating an unfavorable steric

environment for the action of the phospholipase. A situation of steric interference may also be generated by long spacers, i.e. when the number of carbon atoms in the bridging chain is greater than 15.

It is important to note that the functional group of D through which the drug residue is bound to the bridging group may be selected from amino, hydroxyl, thiol and carboxyl groups, with the proviso that when the functional group of D is $-\text{C}(\text{O})\text{OH}$, X is not a carbonyl, and when the functional group of D is $-\text{OH}$, $-\text{NH}_2$, $-\text{NH}$ or $-\text{SH}$, X is a carbonyl group. The above restrictions are applied since some combinations of X with particular drug residues may be unfavorable as yielding a very labile bond which is spontaneously cleaved, therefore greatly lowering the efficacy of the prodrug. Such an unfavorable combination is, for example, where the covalent bond formed is $-(\text{CO})-\text{O}-(\text{CO})-$. This is a labile bond that tends to dissociate, hence greatly reducing drug bioavailability and therapeutic effects at the target site.

The therapeutic efficacy of any particular compound according to the invention should be evaluated by a person skilled in the art considering the general knowledge in chemotherapy and the teachings of the present invention. The choice of a specific compound to be used as a prodrug according to the invention will also depend on the particular disease or disorder to be treated.

In accordance with another aspect of the invention, there are provided pharmaceutical compositions comprising as an active ingredient a prodrug of the general formula I, together with a pharmaceutically acceptable diluent or carrier as are known in the art.

In particularly preferred embodiments of the invention, there are provided pharmaceutical compositions comprising phospholipid derivatives of methotrexate and fluorodeoxyuridine (herein respectively denoted as DP-MTX and DP-FUdR). Specific derivatives are denoted by numerical suffixes.

The linkage of the anti-proliferative drug moiety to the phospholipid is via a bond that is susceptible to cleavage by phospholipase. Thus, at diseased site characterized by elevated activity of phospholipase, DP-MTX prodrug, for example, is cleaved to release free methotrexate or a therapeutically active derivative thereof.

The released drug will inhibit dihydrofolate reductase thus depleting intracellular tetrahydrofolate (FH₄). DNA synthesis and *de novo* purine synthesis will, therefore, also be shut down resulting in inhibition of highly proliferative cells, such as cancer cells. Similarly, at the site of inflammation, the specific release of methotrexate leads to inhibition of leukocyte activation and migration, hence suppressing the inflammatory condition. In the case of rheumatoid arthritis, DP-MTX displays improved intra-articular retention in the inflamed synovium and reduced clearance from the joints due to the presence of its hydrophobic phospholipid moiety.

In a similar way DP-FUdRs, the novel phospholipid derivatives of fluorodeoxyuridine, demonstrate improved properties as anti-proliferative prodrugs. The fact that these prodrugs are capable of selectively releasing their pharmaceutically active component at the target site should enable a significant reduction in therapeutic dose coupled with a reduction in the frequency of administration.

The pharmaceutical compositions of the invention are useful in the treatment of diseases and disorders related to inflammatory, degenerative or atrophic conditions and in the treatment of diseases and disorders related to uncontrolled cell growth. The uncontrolled cell growth may include psoriasis and neoplastic growths including, but not limited to, lymphocytic leukemia, myelogenous leukemia, Burkitt's lymphoma, non-Hodgkin's lymphomas, mycosis fungoides, osteosarcoma, hydatidiform mole, trophoblastic diseases such as chorioadenoma destruens and choriocarcinoma, and carcinomas of the head and neck, breast, liver, lung, colon, ovary, cervix, urinary, bladder, prostate, pancreas, skin, the gastrointestinal tract and the oropharyngeal areas.

The pharmaceutical compositions of the invention are applicable in the treatment of neoplastic growths that may be benign or malignant growths including primary tumors as well as secondary tumors such as metastases.

Furthermore, the pharmaceutical compositions of the invention may be useful for the treatment of drug-resistant tumors, e.g. MTX-resistant and multi-drug resistant (MDR) tumors. This aspect of the invention is based on the unexpected findings that one currently most preferred embodiment of the tested compounds, namely DP-MTX-71, was effective in inhibiting cell growth of drug-resistant tumors.

Moreover, this prodrug is especially advantageous in that it overcomes MTX-resistance without provoking a MDR phenotype.

In another preferred embodiment, the pharmaceutical compositions are useful for treating diseases and disorders related to an inflammatory condition including, but not limited to, granulomatous diseases, arthritis, rheumatoid arthritis, multiple sclerosis, systemic sclerosis, asthma, psoriasis, systemic lupus erythematosus, inflammatory bowel syndromes and migraines.

In yet another preferred embodiment, the pharmaceutical compositions provided by the invention are useful for treating diseases and disorders related to degenerative or atrophic conditions. Said degenerative or atrophic conditions may include, but are not limited to, autoimmune diseases and cerebrovascular and neurodegenerative diseases or disorders in the central and peripheral nervous system.

It should be appreciated that the prodrugs and pharmaceutical compositions in accordance with the invention may be useful in curative as well as in preventive medical treatments. For example, uncontrolled cell growth, inflammatory and degenerative processes possibly leading to pathological conditions and diseases in the nervous system and blood vessels, may be prevented or inhibited by the prodrugs of the invention. In a particular case, the phenomenon of restenosis, frequently developed following invasive procedures used in the treatment of arteriosclerosis diseases, may be ameliorated or prevented by applying a therapeutically effective amount of an anti-proliferative agent in accordance with the invention. The invasive procedures may include, but are not limited to, vascular surgery procedures such as percutaneous transluminal coronary angioplasty (PTCA) and bypass operations.

The pharmaceutical compositions may include therapeutically effective amounts of a prodrug in accordance with the invention together with one or more additional agents known to be effective in the treatment of a particular disease or disorder. For example, beneficial effects have been observed when methotrexate is used as part of a combination therapy in patients with carcinoma or Burkitt's and other non-Hodgkin's lymphomas. Methotrexate is routinely used in combination with cyclophosphamide and fluorouracil in the treatment of breast cancer, in combination with cisplatin and/or doxorubicin in the treatment of cancer of the bladder, and in

combination with cisplatin and bleomycin in the treatment of carcinomas of the cervix and head and neck. In these and other combinations it is possible to substitute one or more of the active drugs for a prodrug according to the invention comprising the corresponding drug residue which may endow the medicament with additional
5 therapeutic value.

The pharmaceutical compositions may be in a liquid, aerosol or solid dosage form, and may be formulated into any suitable formulation including, but not limited to, solutions, suspensions, micelles, emulsions, microemulsions, aerosols, ointments, gels, suppositories, capsules, tablets, and the like, as will be required for the
10 appropriate route of administration.

In yet another aspect, the present invention provides methods for treating pathological conditions related to uncontrolled cell growth, inflammatory and degenerative or atrophic conditions. Said methods comprise administering to an individual in need thereof a therapeutically effective amount of a prodrug of the
15 general formula I or a pharmaceutical composition in accordance with the invention.

The term "therapeutically effective amount" used in the specification refers to the amount of a given prodrug compound according to the invention which antagonizes or inhibits, directly or indirectly, activities associated with inflammatory, degenerative or atrophic processes or uncontrolled cell growth, hence providing either
20 a subjective relief of a symptom(s) or an objectively identifiable improvement as noted by the clinician or other qualified observer.

Any suitable route of administration is encompassed by the invention including, but not being limited to, oral, intravenous, intraperitoneal, intramuscular, subcutaneous, inhalation, intranasal, topical, rectal or other known routes. In preferred
25 embodiments, the pharmaceutical compositions of the invention are orally, intravenously or topically administered.

The dose ranges for administration of the compositions of the invention are those large enough to produce the desired anti-proliferative effect. The dosing range of the prodrug varies with the specific drug used, the treated pathological condition, the
30 route of administration and the potency of the particular prodrug molecule in releasing the drug at the specific target site. The dosage administered will be dependent upon the

age, sex, health, weight of the recipient, concurrent treatment, if any, frequency of treatment and the nature of the effect desired. Dosage regimen and means of administration will be determined by the attending physician or other person skilled in the art.

5

The invention will now be illustrated by the following non-limiting examples, which include detailed description of some specific embodiments of the invention. These examples are illustrative and are not to be read as limiting the scope of the invention as is defined by the claims which follow.

10

EXAMPLES

I. CHEMICAL EXAMPLES

Example 1: General synthesis of phospholipid derivatives of methotrexate (DP-MTX)

15

A general procedure for the synthesis of DP-MTX compounds is described below. It is important to note that in this protocol a selective binding to the α and γ positions of the methotrexate has been achieved.

Binding between the phospholipid and the MTX in these derivatives is via an ω -aminoacid at position-2 of lipid moiety. The synthesis pathway consists of six steps. Step 1 is protection of an amino group with a benzyloxycarbonyl resulting in the corresponding Z-amino acid. In step 2 the Z-amino acid is converted into the Z amino acid anhydride. Step 3 is synthesis of DP-amino acid, namely the formation of lipid derivative comprising the amino acid with protected amino group and a lyso-
lecithin. Step 4 is de-protection, by hydrogenation, of the amino group of the amino acid. In step 5 the mixed anhydride of methotrexate or its γ -esters are prepared by reacting methotrexate (γ -esters) with isobutyl chloroformate in the presence of triethyl amine. In the last stage, step 6, the final product is obtained by reacting the mixed anhydride of methotrexate, or its ester, with 1-acyl-2-(ω -amino) acyl-Sn-glycero-3-phosphatidylcholine in the presence of triethylamine as a catalyst.

20

25

30

For the sake of clarity, each of the intermediary steps will be described below in details.

Stage 1. Protection of the amino group of amino acid (Preparation of Z-amino acid).

5 To a mixture of 0.1 mol corresponding amino acid (3-aminopropanoic acid, 4-aminobutanoic acid, 6-aminohexanoic acid, 8-aminooctanoic acid etc.) in ethanol (25 ml) in round-bottom flask (250 ml) equipped with a magnetic stirrer and dropped funnel, a solution of NaOH (8.8g., 0.22 mol) in 100 ml water is added and the mixture is stirred by magnetic stirrer until fully dissolved. The obtained solution is cooled to
10 0°C in an ice-water bath, and benzyl chloroformate (27.4g , 0.15 mol) is added drop wise over 30 min. The reaction mixture is stirred for 3 hours at 0°C. Subsequently, about 100 ml water is added to the reaction solution and the mixture is poured into separated funnel. The water solution is extracted with diethyl ether (3X 50 ml). The water phase is separated and acidified with HCl (3N) to pH=1 while cooling in an ice-
15 water bath. If a precipitate is formed, it is filtered, washed with water and dissolved in 100 ml chloroform. The chloroform solution is dried with sodium sulfate for two hours. Then the sodium sulfate is separated from the chloroform solution by filtration and the solvent is evaporated in evaporator under vacuum. The residue is washed with hexane, and dried overnight in vacuum over phosphorus pentoxide (P₂O₅).

20 If a precipitate is not formed, or in order to maximize the product yield, the acidified aqueous fraction is washed with chloroform (2X 50 ml). The chloroform extracts are combined and washed with water (50ml). The following operations with this solution are the same as for the above-described chloroform solution of the precipitate, namely, drying with sodium sulfate for two hours, then separating the
25 sodium sulfate from the chloroform solution by filtration and evaporating the solvent in evaporator under vacuum. The residue is then washed with hexane, and dried overnight in vacuum over phosphorus pentoxide.

30 All products were analyzed on TLC as follows: TLC analysis. Silica gel 60 on aluminium sheet. Eluent is chloroform-methanol (4:1 v/v). Indicator is a spray of the composition: 4-methoxybenzaldehyde (10ml), absolute ethanol (200ml), 98% sulfuric

acid (10ml) and glacial acetic acid (2ml). The chromatogram is sprayed with the indicator and then charred at 150-180°C.

The following are specific intermediate products obtained at the end of stage 1 of the synthesis procedure:

- 5 3-[N-(Benzyloxycarbonyl)amino]propanoic acid. $C_6H_5-CH_2-O-C(O)-NH-CH_2-CH_2-COOH$.

White solid. Yield 60%. TLC analysis: One spot R_f 0.7.

1H NMR (CD_3OD), $\delta(ppm)$: 2.46-2.52 (t, 2H, $\alpha-CH_2$ group of alanine), 3.29-3.39 (t, 2H, $\beta-CH_2$ group of alanine) 5.06 (s, 2H, benzylic CH_2 group), 7.27-7.32 (m, 5H, C_6H_5 group).

10

- 4-[N-(Benzyloxycarbonyl)amino]butanoic acid. $C_6H_5CH_2-O-C(O)-NH-(CH_2)_3-COOH$.

White solid. Yield 60%. TLC analysis: One spot. R_f 0.7.

1H NMR (CD_3OD), $\delta(ppm)$: 1.71-1.82 (m, 2H), 2.28-2.34 (m, 2H), 3.10-3.17 (t, 2H), 5.06 (s, 2H), 7.26-7.34 (m, 5H).

15

- 5-[N-(Benzyloxycarbonyl)amino]valeric acid. $C_6H_5CH_2-O-C(O)-NH-(CH_2)_4-COOH$.

White solid. Yield 60%. TLC analysis: One spot. R_f =0.7.

1H NMR (CD_3OD), $\delta(ppm)$: 1.45-1.50 (m, 2H), 1.56-1.62 (m, 2H), 2.25-2.31 (t, 2H), 3.08-3.13 (t, 2H), 5.05 (s, 2H), 7.26-7.34 (m, 5H).

- 20 6-[N-(Benzyloxycarbonyl)amino]hexanoic acid. $C_6H_5CH_2-O-C(O)-NH-(CH_2)_5-COOH$.

White solid. Yield 50%. TLC analysis: One spot. R_f 0.7.

1H NMR (CD_3OD), $\delta(ppm)$: 1.30-1.63 (m, 6H), 2.24-2.30 (t, 2H), 3.07-3.13 (t, 2H), 5.05 (s, 2H), 7.29-7.34 (m, 5H).

- 25 8-[N-(Benzyloxycarbonyl)amino]octanoic acid. $C_6H_5CH_2-O-C(O)-NH-(CH_2)_7-COOH$.

White solid. Yield 50%. TLC analysis: One spot. R_f 0.7.

1H NMR (CD_3OD), $\delta(ppm)$: 1.32 (bs, 6H), 1.47-1.50 (m, 2H), 1.53-1.59 (m, 2H), 2.23-2.29 (t, 2H), 3.06-3.12 (t, 2H), 5.05 (s, 2H), 7.29-7.34 (m, 5H).

Stage 2. Synthesis of Z-amino acid anhydride.

- 30 The solution of the corresponding Z-amino acid produced at stage 1 (0.05 mol) in freshly distilled dichloromethane (25 ml) is introduced, under an atmosphere of

argon, into double-neck round-bottom equipped with magnetic stirrer and dropped funnel. A solution of dicyclohexylcarbodiimide (DCC) (0.0325 mol) in 25 ml of freshly distilled dichloromethane, also under argon, is added drop wise, with stirring, to the solution of Z-amino acid. After 20 min of stirring, the obtained precipitate of urea is filtered and the solution evaporated under vacuum. The crude residue is washed with hexane (2X 20 ml) to remove remaining DCC and then dried in vacuum.

TLC analysis: The same procedure is used for TLC analysis of anhydride of the all Z-amino-acids. Silica gel 60 on aluminium sheet. Eluent is the mixture of chloroform with methanol (9:1 v/v). For indication, ninhydrine spray is used on the chromatogram followed by charring (100-150°C).

The following are specific intermediate products obtained at the end of stage 2:

Anhydride of Z-(3-amino)propanoic acid. White solid. Yield is 70%.

TLC analysis: One spot R_f 0.8.

Anhydride of Z-(4-amino)butanoic acid. White solid. Yield is 70%.

TLC analysis: One spot. R_f 0.8.

Anhydride of Z-(5-amino) valeric acid. White solid. Yield is 70%.

TLC analysis: One spot. R_f 0.8.

Anhydride of Z-(6-amino)hexanoic acid. White solid. Yield is 70%.

TLC analysis: One spot. R_f 0.8.

Anhydride of Z-(8-amino)octanoic acid. White solid. Yield is 75%.

TLC analysis: One spot. R_f 0.85.

Stage 3. Preparation of 1-acyl-2- (Z-amino)acyl-sn-glycero-3-phosphotidylcholine.

The anhydride of the corresponding Z-amino acid, 0.01 mol dissolved in 150 ml of freshly distilled chloroform, is introduced, under an inert atmosphere of argon, into a single-neck round-bottom flask (250 ml) equipped with a magnetic stirrer. To this solution 0.01 mol (1.22 g) 4-(dimethylamino)pyridine (DMAP) in 25 ml chloroform is added, followed by addition of a suspension of 0.0056 moles lyso-lecithin in 50 ml of chloroform. The reaction mixture is vigorously stirred for 3-5 hours at room temperature. The lyso-lecithin dissolves and reaction mixture becomes transparent after about 2 hours of stirring. The reaction is monitored by TLC using

silica gel 60 on aluminium sheet, the eluent is chloroform:methanol:water, 65:35:5, the indicator is a spray of the composition: 4-methoxybenzaldehyde (10 ml), absolute ethanol (200 ml), 98 % sulfuric acid (10 ml) and glacial acetic acid (2 ml). The chromatogram is sprayed with the indicator followed by charring at 150°C. The reaction is assumed to be complete and stopped when all the lyso-lecithin has disappeared. The reaction mixture is then transferred into a separating funnel and washed with a solution of 1% HCl (3x 50 ml), then with saturated solution of sodium bicarbonate (3x 50 ml) and finally with water (3x 50 ml). The obtained product in the organic solution is dried over sodium sulfate and then filtered. The solvent is

evaporated at 30°C in vacuo and the residue is washed with hexane and left to dry overnight under vacuum. The resulted molecule 1-acyl-2-(Z-amino)acyl-sn-glycero-3-phosphatidylcholine is the main product of the reaction.

TLC analysis: Silica gel 60 on aluminium sheet. Eluent is chloroform/methanol/water (65:35:5 v/v). Indicator is a spray of the composition: 4-methoxybenzaldehyde (10 ml), absolute ethanol (200 ml), 98% sulfuric acid (10 ml) and glacial acetic acid (2 ml). The chromatogram is sprayed with the indicator and then charred at 100-150°C. The following are specific intermediate products obtained at the end of stage 3:

1-Stearoyl-2-{3'-[N-(Benzyloxycarbonyl)amino]}propanoyl-sn-glycero-3-phosphatidyl choline.

White wax. Yield 70%. TLC analysis: One spot. R_f 0.55

1-Stearoyl-2-{4'-[N-(Benzyloxycarbonyl)amino]}butanoyl-sn-glycero-3-phosphatidyl choline.

White wax. Yield 70%. TLC analysis: One spot. R_f 0.55.

1-Stearoyl-2-{5'-[N-(Benzyloxycarbonyl)amino]}valeroyl-sn-glycero-3-phosphatidyl choline.

White wax. Yield 65%. TLC analysis: One spot. R_f 0.55.

1-Stearoyl-2-{6'-[N-(Benzyloxycarbonyl)amino]}hexanoyl-sn-glycero-3-phosphatidyl choline.

White wax. Yield 65%. TLC analysis: One spot. R_f 0.55.

1-Stearoyl-2-{8'-[N-(Benzyloxycarbonyl)amino]}octanoyl-sn-glycero-3-phosphatidyl choline.

White wax. Yield 65 %. TLC analysis: One spot. R_f 0.55.

Stage 4. Reduction of amide group of amino acyl lipid.

The obtained 1-stearoyl-2- $\{\omega$ -[(N-Benzyloxycarbonyl)amino]}acyl-3-phosphatidyl-choline (0.0025 mol) is dissolved in a mixture of 100 ml methanol and 5 ml acetic acid. The solution is introduced into round bottom double neck flask (200 ml) equipped with a magnetic stirrer, under an atmosphere of argon. Pd/C (0.5 g) is added to the solution and hydrogen is blown through the reaction mixture for 4 hours. The reaction proceeding is monitored by TLC analysis under the following conditions: silica gel 60 on aluminium sheet, eluent is the mixture of chloroform/methanol/water (65:35:5 v/v), indicator is a spray of the composition: p-methoxybenzaldehyde (10 ml), absolute ethanol (200 ml), 98% sulfuric acid (10 ml) and glacial acetic acid (2 ml). The chromatogram is sprayed with the indicator and then charred at 100-150°C.

The reaction assumed to be complete and hydrogenation is stopped after the corresponding 1-stearoyl-2-Benzyloxycarbonylaminoacyl-sn-glycero-phosphatidylcholine has disappeared from the reaction mixture. The reaction mixture is then filtered through Wattman paper to remove the Pd/C, evaporated at 30°C, under vacuum. The crude residue is washed with ether (3x 30 ml) and dried in vacuo overnight.

Conditions of the TLC analysis are the same as indicated above. The following are specific intermediate products obtained at the end of stage 4:

1-Stearoyl-2-(3-amino)propanoyl-sn-glycero-3-phosphatidylcholine, acetic acid.

White wax. Yield 70%. TLC analysis: One spot. R_f 0.2.

1-Stearoyl-2-(4-amino)butanoyl-sn-glycero-3-phosphatidylcholine, acetic acid.

White wax. Yield 70%. TLC analysis: One spot. R_f 0.2.

1-Stearoyl-2-(5-amino)valeroyl-sn-glycero-3-phosphatidylcholine, acetic acid.

White wax. Yield 65%. TLC analysis: One spot. R_f 0.2.

1-Stearoyl-2-(6-amino)hexanoyl-sn-glycero-3-phosphatidylcholine, acetic acid.

White wax. Yield 65%. TLC analysis: One spot. R_f 0.2.

1-Stearoyl-2-(8-amino)octanoyl-sn-glycero-3-phosphatidylcholine, acetic acid.

White wax. Yield 65 / . TLC analysis: One spot. R_f 0.2.

Stage 5. Synthesis of the mixed anhydride of methotrexate.

Freshly distilled dimethylformamide (DMF) (25 ml) and THF (75 ml) is introduced, under atmosphere of argon, into a double-neck round-bottom flask (150 ml), equipped by magnetic stirrer and thermometer. Methotrexate (MTX; 0.454g., 1 mmol) and triethylamine (0.182g., 239 μ l, 1.8 mmol) are added, under the same conditions, and dissolved in the solvent with stirring.

In protocols for the synthesis of DP-ester-MTX compounds, the MTX was replaced by a α - or γ -ester derivative of MTX wherein the hydrogen atom of the α or γ carboxylic group of methotrexate is substituted by an alkyl group of from 2 to 30 carbon atoms. The ester derivatives of MTX were prepared according the procedure described by Rosowsky et al. (J. Medicin. Chem., 1978, 21: 380-386). The reaction mixture is then cooled to -25°C in dry ice-acetone bath. Isobutyl chloroformate (0.166 g., 163 μ l, 1.2 mmol) is added dropwise to the reaction mixture over 30 min. During this procedure and the following 30-min, the temperature of the reaction mixture is maintained at -25°C . The obtained product is a white precipitate, which is not extracted from reaction mixture but was directly used in the next stage of the synthesis.

Stage 6. Synthesis of α -or γ -monoamide of methotrexate (DP- α -MTX monoamide and DP- γ -MTX monoamide, respectively)

A solution comprising the corresponding 1-stearoyl-2- ω -aminoacyl-sn-glycero-3-phosphatidylcholine of stage 4, acetic acid (0.589g., 0.9 mmol) and triethylamine (0.182g., 239 μ l, 1.8 mmol) in 30 ml of dry freshly distilled chloroform is added dropwise to the reaction mixture of stage 5 for 30 min at -25°C . The reaction mixture is stirred for additional one hour at -25°C , and then for overnight at room temperature. The solvents are removed in evaporator under reduced pressure. The obtained residue is a thick viscous liquid. Diethyl ether (50 ml) is added to this liquid and the mixture is stirred. The product is gradually transformed into a yellow powder which is filtered and washed with diethyl ether (3x 20 ml). The crude product is purified by column chromatography as follows: 450 g. crude product is dissolved in 50 ml of methanol, followed by addition of 11.0 g. dry silica gel. The mixture is

swirled, and then the volatile liquid is evaporated under reduced pressure to yield free-flowing yellow finely divided solid. The obtained solid is packed on top of a silica column (3x30 cm) (100g. silica gel per 1g of crude product). The product is eluted in succession with the solution MeOH:H₂O of variable composition: first
5 fraction is 100:0 (v/v) (1L), second fraction is 99:1 (v/v) (1L), third fraction is 98:2 (v/v) (1L), and fourth fraction is 98:3 (v/v) (2L). The fractions, which contain the product (the determination is carried out by TLC analysis), are combined and the solvent is evaporated under reduced pressure. The obtained product (about 250 mg) is dissolved in mixture of methanol (10 ml) and chloroform (250 ml). The solution is
10 washed with 1% HCl (3x 20 ml) and then with water (3x 20 ml). To achieve better separation of the aqueous and organic phases, isopropanol (about 25% of the volume of solution) is added. Isopropanol addition also promotes transition of the product into the organic phase. The organic layers are combined and dried over sodium sulfate. The sodium sulfate is filtered off and the solvent is distilled under reduced
15 pressure. The obtained product is dried under vacuum for 3 hours.

Examples of resulted final products and their analyses are:

1-Stearoyl-2-[3-(α -MTX amido)]-propanoyl-sn-glycero-3-phosphatidylcholine.
C₄₉H₇₉N₁₀O₁₂P.

Yield 35%. Yellow solid. Decompose at 200°C without melting. pH 5.1.

20 TLC analysis: Silica gel on aluminium plates. Eluent is CH₃OH:H₂O (98:2, v/v). Indication by UV-Vis spectrum. One spot. R_f 0.16.

MS(FAB). C₄₉H₇₉N₁₀O₁₂P. Main peak (+FAB) is 1031.2.

¹H NMR (CD₃OD), δ (ppm): 0.89 (t, 3H), 1.22-1.29 (broad s, 30H), 1.53 (m, 2H), 2.08 (m, 2H), 2.35 (m, 2H), 2.60 (t, 2H), 2.50 (t, 2H), 3.21 (s, 3H), 3.38 (s, 9H), 3.47
25 (t, 2H), 3.93-4.31 (m, 8H), 4.41 (m, 1H), 4.80 (s, 2H), 5.21 (m, 1H), 6.84 (d, 2H), 7.76 (d, 2H), 8.60 (s, 1H).

³¹P NMR (CD₃OD), δ (ppm): -3.3 (s).

Chemical analysis: C₄₉H₇₉N₁₀O₁₂P.HCl.3H₂O. Calculated: C 52.47%, H 7.55%, N 12.49%, P 2.76%, Cl 3.16%. Found: C 52.92%, H 7.76%, N 12.21%, P 2.46%, Cl
30 3.08%.

1-Stearyl-2-[6-(α -MTX amido)]-hexanoyl-sn-glycero-3-phosphatidylcholineC₅₂H₈₅N₁₀O₁₂P.

Yield 40 %. Yellow solid. Decomposes without melting at 200°C. pH 5.0.

TLC analysis: Silica gel on aluminium plates. Eluent is MeOH:H₂O (98:2, v/v).5 Indication is UV-Vis spectra. One spot R_f 0.18.MS (FAB): C₅₂H₈₅N₁₀O₁₂P 1073.3 (+FAB) is main peak.¹H NMR (CD₃OD), δ (ppm): 0.86-0.89 (t, 3H), 1.22-1.30 (broad s, 32H), 1.51-1.55 (m, 4H), 2.06-2.10 (m, 4H), 2.33-2.36(m, 2H), 2.58-2.62 (t, 2H), 2.49-2.52 (t, 2H), 3.22 (s, 3H), 3.38 (s, 9H), 3.45-3.48 (t, 2H), 3.92-4.35 (several m, 8H), 4.40-4.42 (m, 1H), 4.79 (s, 2H), 5.20 (m, 1H), 6.82-6.84 (d, 2H), 7.74-7.76 (d, 2H), 8.58 (s, 1H).³¹P NMR (CD₃OD), δ (ppm): -3.7 (s).Chemical analysis: C₅₂H₈₅N₁₀O₁₂P.2HCl.5H₂O: Calculated: C 50.48%, H 7.84%, N 11.33%, P 2.51%, Cl 5.74%. Found: C 50.45%, H 7.22%, N 11.46%, P 2.43%, Cl 5.43%.

15

1-Stearyl-2-[3-(α - dodecylate- γ -MTX amido)]-propanoyl-sn-glycero-3-phosphatidylcholine. C₆₁H₁₀₃N₁₀O₁₂P.

Yield 40%. Yellow solid. Decompose at 200°C without melting.

TLC analysis: Silica gel on aluminium plates. Eluent is CHCl₃:CH₃OH:H₂O (65:35:5, v/v). Indication by UV-Vis spectrum. One spot. R_f 0.17.MS(FAB). C₆₁H₁₀₃N₁₀O₁₂P. Main peak (+FAB) is 1198.4.¹H NMR (CD₃OD + CDCl₃), δ (ppm): 0.82-0.85 (t, 3H), 1.23 (broad s, 46H), 1.53-1.58 (m, 2H), 2.24-2.34 (m, 3H), 2.50-2.52 (m, H), 3.17 (s, 9H), 3.41 (t, 1H), 3.54 (s, 1H), 4.04-4.18 (m, 4H), 4.32-4.47 (m, 1H), 4.80 (s, 2H), 5.21 (m, 1H), 6.79-6.82 (d, 2H), 7.76 (d, 2H), 8.54 (s, 1H).³¹P NMR (CD₃OD + CDCl₃), δ (ppm): +0.71 (s).Chemical analysis: C₆₁H₁₀₃N₁₀O₁₂P.3H₂O. Calculated: C 58.47%, H 8.71%, N 11.18%, P 2.47% . Found: C 58.67%, H 8.88%, N 11.15%, P 2.47%.

30

Example 2: General synthesis of phospholipid ester derivatives of methotrexate (DP-MTX esters)

The synthetic procedure for the preparation of DP-MTX-esters is a three-stage process.

5 **Stage 1. Synthesis of ω -bromoalkylcarboxylic anhydride.**

The solution of the corresponding ω -bromoalkyl acid (0.05 mol) in freshly distilled dichloromethane (25 ml) is introduced, under an inert atmosphere of argon, into double-neck round-bottom equipped with magnetic stirrer and dropping funnel. A solution of dicyclohexylcarbodiimide (DCC) (0.0325 mol) in 25 ml freshly distilled
10 dichloromethane, also under argon, is added drop wise, with stirring, to the solution of Z-amino acid. After 20 min. of stirring, the produced urea precipitate is filtered and the obtained solution is evaporated under vacuum. The crude residue is washed with hexane (2x 20 ml) to remove remaining DCC and then dried in vacuum.

15 **Stage 2. Preparation of 1-acyl-2- ω -bromoalkylcarboxy-sn-glycero-3-phosphatidylcholine.**

0.01 mol of the corresponding ω -bromoalkyl-carboxylic anhydride (obtained in stage 1) dissolved in 150 ml freshly distilled chloroform, is introduced, under an inert atmosphere of argon, into a single-neck round-bottom flask (250 ml) equipped with a magnetic stirrer. To this solution 0.01 mol (1.22 g) 4-(dimethylamino)pyridine
20 (DMAP) in 25 ml chloroform is added, followed by addition of a suspension of 0.0056 moles lyso-lecithin in 50 ml chloroform. The reaction mixture is vigorously stirred for 3-5 hours at room temperature. The lyso-lecithin dissolves and reaction mixture becomes transparent after about 2 hours of stirring. The reaction is monitored by TLC using silica gel 60 on aluminium sheet, the eluent is
25 chloroform:methanol:water, 65:35:5, the indicator is a spray of the composition: 4-methoxybenzaldehyde (10 ml), absolute ethanol (200 ml), 98% sulfuric acid (10 ml) and glacial acetic acid (2 ml). The chromatogram is sprayed with the indicator followed by charring with hot air at 150°C. The reaction is assumed to be complete and is stopped when all the lyso-lecithin has disappeared. The reaction mixture is then
30 transferred into a separating funnel and washed with a solution of 1 / HCl (3x 50 ml), then with saturated solution of sodium bicarbonate (3x 50 ml) and finally with water

(3x 50 ml). The organic phase is dried over sodium sulfate and then filtered. The solvent is evaporated at 30°C in vacuo and the obtained residue is washed with hexane and left to dry overnight under vacuum. The desired molecule 1-acyl-2- ω -bromoalkylcarboxy-sn-glycero-3-phosphatidylcholine is the main product of the reaction.

Stage 3. Synthesis of α -monoesters of methotrexate (DP- α -MTX esters)

A solution of 1-Stearoyl-2- ω -bromoalkylcarboxy-sn-glycero-3-phosphocholine (0.016 mole) in 50 ml dimethyl sulfoxide (DMSO) (freshly distilled over CaH₂) is added to a mixture of methotrexate free acid (0.008 mole) and CsCO₃ (0.008 mole) in DMSO (150 ml) under an atmosphere of argon. The reaction mixture is stirred at room temperature for 48 hours. The solvent is evaporated under vacuum (1-2 mm Hg, 50°C). The obtained residue is acidified with 1% HCl to pH=3, followed by an overnight incubation at 0°C. The mixture is then filtered, washed with water and dried over P₂O₅ under vacuum. The resultant product is 1-stearoyl-2-[ω -(α -MTX)alkyl]-oxycarbonyl-sn-glycero-3-phosphocholine. This crude product is purified by column chromatography (1 g. crude product per 100 g. Silica gel 60) using the eluent: CHCl₃ : MeOH : H₂O (v/v 40:9:1)

Example 3: Synthesis of phospholipid derivatives of 2'-deoxy-5-fluorouridine (DP-5FUdR)

The procedure for the synthesis of DP-5FUdR compounds of the invention is exemplified below by the synthesis of the specific compound 1-stearoyl-2-[5'-(2'-deoxy-5'-fluorouridine-5'-)-3'',3''-dimethyl]glutaroyl-1''-sn-glycero-3-phosphatidylcholine. The detailed description of the synthesis is described below. The products of the different stages of the synthesis are denoted as compounds (1) to (5).

Stage 1. Preparation of 5'-O-Trityl 5-fluoro-2'-deoxyuridine (1).

5-Fluoro-2'-deoxyuridine (2 g, 8.8 mmol) was dissolved in 40 ml dry pyridine under an atmosphere of nitrogen. Trityl chloride (3.6 g, 12.6 mmol) was added and the reaction mixture stirred at room temperature for 48 hours. The reaction was quenched

with methanol (20 ml), and then the mixture was concentrated to dryness under reduced pressure. The compound was purified on silica gel (flash), using chloroform/methanol (97/3) as an eluent. 4.1 g of the product (1) was obtained as a colorless, crystalline solid. Yield – 99%.

5 **Stage 2. Preparation of 3'-O-Levulinoyl 5-fluoro-2'-deoxyuridine (2).**

The mixture of levulinic acid (2.32 g, 20 mmol), N,N'-dicyclohexylcarbodiimid (2.06 g, 10 mmol) and ether (60 ml) was stirred at room temperature for 5 hours, followed by filtration and evaporation of the solvent. The obtained compound was dissolved in anhydrous pyridine (10 ml). To the solution, compound (1) (4 g, 8.5
10 mmol) was added. The mixture was stirred for 24 hours at room temperature. Then ~ 50 g chipped ice was added and the reaction mixture was stirred until all the ice melted. The reaction mixture was extracted with chloroform (4 x 20 ml), and the combined extracts were dried over anhydrous MgSO₄. Concentration under reduced pressure followed by removal of excess pyridine under high vacuum yielded 3'-
15 levulinoyl-5'-trityl-5-fluoro-2'-deoxyuridine as yellow oil. This product was re-dissolved in 80% acetic acid (20 ml) and heated for 20 min at 100 °C. The reaction mixture was then evaporated under reduced pressure, followed by repeated evaporation from benzene in order to remove traces of water and acetic acid. The resultant reaction mixture was then purified by flash chromatography on silica gel,
20 using chloroform/methanol (40/1). 1.3 g of compound (2) was obtained as white powdery solid. Yield - 46%.

Stage 3. Preparation of 1-Stearoyl-2- 3'',3''-dimethyl-glutaroyl-sn-glycero-3-phosphocholine (3).

A solution of 1-stearoyl-sn-glycero-3-phosphocholine (1.8 g, 3.6 mmol), 3,3-
25 dimethylglutaric anhydride (3.5 g, 25.2 mmol), sodium salt of 2-propylpentanoic acid (0.5 g, 3.2 mmol) in N,N-dimethylformamide (20 ml) was heated for 5 hours at 90 °C. The reaction mixture was concentrated to dryness under high vacuum. The residue was dissolved in chloroform/methanol (2/1) 100 ml and washed with 0.1 N HCl (2 x 50 ml). After evaporation and flash chromatography on silica gel (CHCl₃/MeOH/H₂O/AcOH
30 60/30/4/1), 2 g of compound (3) was obtained as white crystalline solid. Yield – 87 / .

Stage 4. Preparation of 1-Stearoyl-2-[5''-O-(-3'-O-levulinyl-5-fluoro-2'-deoxyuridine)-3'',3''-dimethylglutaroyl]-sn-glycero-3-phosphocholine (4).

To 1 g of compound (3) (1.5 mmol), 2.0M solution of oxalyl chloride in dichloromethane (20 ml, 4 mmol) was added. The mixture was stirred overnight at 20 °C. The acid chloride was separated from the oxalyl chloride by removal of the volatiles in vacuum followed by two cycles of dissolution in dry benzene (20 ml) and evaporation.

A solution consisting of compound (2) (700 mg, 2.15 mmol) and triethylamine (0.22 g, 2.17 mmol) in dichloromethane (10 ml) was added drop wise to solution of the acid chloride in dichloromethane (10 ml) at 0 °C. The mixture was left overnight at 20 °C, then the volatile solvents were removed and the residue was dissolved in chloroform/methanol (2/1, 50 ml) and washed with 0.1N HCl (20 ml). The solvent was removed and the residue was subjected to chromatography on silica gel using methanol/chloroform (10-33%) followed by water/methanol/chloroform (2/30/60) to yield 700 mg of compound (4).

Yield - 48%.

Stage 5. Preparation of 1-stearoyl-2-[5'''-(2''-deoxy-5'-fluorouridine-5''-)-3''',3'''-dimethyl]glutaroyl-1'''-sn-glycero-3-phosphatidylcholine (5).

Hydrazine monohydrate (0.25 ml, 4.8 mmol) in pyridine/acetic acid (3:2 v/v; 5 ml) was added to compound (4) (200 mg, 0.2 mmol) dissolved in pyridine (5 ml). After 2 min at 20 °C, the solution was cooled to 0 °C and 2-pentanone (2 ml) was added. The reaction mixture was then evaporated to dryness. Preparative TLC using water/methanol/chloroform (5/35/60) as the solvent yielded 120 mg of the final product (5). Yield - 68%.

II. BIOLOGICAL EXAMPLES:

Example 4: Evaluation of anti-proliferative activity of methotrexate derivatives (in vitro study)

A series of methotrexate derivatives were screened for their inhibitory effect on cell growth. Three tumor cell lines were used for this screen: i) human leukemia cell line CCRF-CEM which is MTX-sensitive (herein denoted as CEM), ii) a mutant of CEM cell line which is MTX-resistant due to a mutation in the RFC (herein denoted as CEM-MTX; provided by Dr. Y. Assraf, Technion, Israel), and iii) rat C6 glioma cell line.

Cells were seeded at a density of 10^4 cells/ml in RPMI medium supplemented with 10% FCS in 96 well plates. The cultured cells were incubated, during their linear growth phase, in the presence and absence of various concentrations of MTX derivatives. The tested compounds were prepared as 1 mM stock in absolute ethanol or DMSO and were diluted into the medium. The highest vehicle presence in the assay was 5% of the vehicle. After 72 hours at 37°C the cytotoxic effect on the cells was estimated by using the colorimetric MTT assay (Mosmann (1983) J. Immunol Methods 65: 55-63) that measures mitochondrial reductase activity and serves for quantitative assessment of cellular viability. Drug concentration required for inhibiting the growth of 50% of cell population in culture over a 72 hour period incubation, is defined as EC_{50} . The EC_{50} values were calculated from dose response curves for each tested compound. The results of this screen are presented in Table 1. The following MTX derivatives were tested:

α - dodecylate-MTX (MTX-47);

γ -dodecylate-MTX (MTX-48);

α - dodecylate-MTX- γ -dodecylate (MTX-256);

1-Stearoyl-2-[3-(α -dodecylate- γ -MTX amido)-Propanoyl]-sn-Glycero-3-Phosphatidylcholine (denoted DP-MTX-71);

1-Stearoyl-2-[3-(α -MTX amido)-Propanoyl]-sn-Glycero-3-Phosphatidylcholine (denoted DP-MTX-93);

1-Stearoyl-2-[4-(α -MTX amido)-Butanoyl]-sn-Glycero-3- Phosphatidylcholine (denoted DP-MTX-129);

1-Stearoyl-2-[6-(α -MTX amido)-Hexanoyl]-sn-Glycero-3- Phosphatidylcholine (denoted **DP-MTX-106**);

1-Stearoyl-2-[8-(α -MTX amido)-Octanoyl]-sn-Glycero-3- Phosphatidylcholine (denoted **DP-MTX-142**);

5 1-Stearoyl-2-[3-(γ -dodecylate- α -MTX amido)-Propanoyl]-sn-Glycero-3- Phosphatidylcholine (denoted **DP-MTX-128**); and

1-Stearoyl-2-[8-(γ -dodecylate- α -MTX amido)-Octanoyl]-sn-Glycero-3- Phosphatidylcholine (denoted **DP-MTX-127**).

The cytotoxic activity of the DP-MTX derivatives on the tested cells was compared to
10 that of commercial MTX (Abitrexate®, from ABIC, Israel; or Sigma cat. # M 8407).

Table 1: Effect of various derivatives of MTX on cell growth

Compound	EC ₅₀ , nMol CEM	EC ₅₀ , nMol CEM-MTX	EC ₅₀ , nMol C6 glioma
MTX	21	2535	46
α -C ₁₂ H ₂₅ -MTX	88	9423	
γ -C ₁₂ H ₂₅ -MTX	13	4864	
α -C ₁₂ H ₂₅ -MTX- γ -C ₁₂ H ₂₅	4855	5093	807
DP-MTX-71	169	213	14
DP-MTX-93	27	4651	417
DP-MTX-129	821	5982	3109
DP-MTX-106	499	8398	3219
DP-MTX-142	6344	11591	35982
DP-MTX-128	771	1602	
DP-MTX-127			795

15

As can be seen from the results in Table 1, the various tested DP-MTX derivatives were active to a different degree in their ability to inhibit cell growth of the human leukemia cell lines and the rat glioma cells. Among the tested DP-MTX

derivatives, the most effective in inhibiting CEM cell growth was DP-MTX-93, having EC_{50} value which is similar to that of the parent drug MTX. Another compound, DP-MTX-71, was found to exhibit a significantly improved effect in inhibiting cell growth of the resistant cell line CEM-MTX. The DP-MTX-71 derivative of MTX was about 10 fold more active in its cell growth inhibition effect on the resistant cell line in comparison to the parent drug, MTX.

DP-MTX-71 has demonstrated similar EC_{50} values for both the leukemia cell line CEM and the mutant MTX-resistant cell line, CEM-MTX and was found to be especially effective in inhibiting growth of the C6 glioma cell line. It is important to note that α - $C_{12}H_{25}$ -MTX, γ - $C_{12}H_{25}$ -MTX and the di-ester α - $C_{12}H_{25}$ -MTX- γ - $C_{12}H_{25}$ were not effective in inhibiting the growth of the MTX-resistant cell line, CEM-MTX.

Conclusion: DP-MTX-93 and DP-MTX-71 were identified as the most efficacious of the tested MTX-derivative compounds. The DP-MTX-71 compound that, unexpectedly, was found to be active also in inhibiting the MTX-resistant tumor cell line, was chosen to be used in further experiments aimed to analyze in more detail the cytotoxic effect of DP-MTX.

Example 5: Evaluation of cytotoxicity of methotrexate derivatives (in vitro)

The toxicity of methotrexate derivatives of the invention (DP-MTX) was evaluated in an additional cell-culture system. The doses required to inhibit the growth of cultured Friend cells (an erythroleukemic cell line) by 50%, over a 72 hour period (ED_{50}), were determined using a mitochondrial viability (MTT) endpoint assay.

Friend cells were seeded at a density of 10^4 cells/ml in RPMI medium supplemented with 10% FCS in 96 well plates. The cultured cells were incubated, during their linear growth phase, in the presence and absence of various concentrations of MTX derivatives. After 72 hours at $37^{\circ}C$, cell viability was estimated by using the colorimetric MTT assay.

Table 2

Compound	n	R	ED ₅₀ , nMol
MTX (control)	-	-	10
<u>α-MTX</u>			
DP-MTX-93	2	H	97.5
DP-MTX-129	3	H	97.5
DP-MTX-106	5	H	390
DP-MTX-142	7	H	3000
<u>γ-MTX</u>			
DP-MTX-71	2	C ₁₂ H ₂₅	23

n is the number of carbon atoms in the linker between the MTX residue and the phospholipid moiety; R is the substituent at the γ - or α -carboxylic group of the α -MTX and γ -MTX compounds, respectively.

Of the tested MTX derivatives, DP-MTX-93, DP-MTX-129, DP-MTX-106 and DP-MTX-142 are α -MTX derivatives, namely molecules wherein MTX is linked to the lipid moiety through its α -carboxylic group. MTX-71, on the other hand, is a γ -MTX derivative, namely a molecule wherein MTX is linked through its γ -carboxyl instead of its α -carboxylic group. In addition, MTX-71 has an R residue, C₁₂H₂₅ group, linked to its α -carboxyl.

As can be seen in Table 2, DP-MTX-71 was about 2-fold less active than the parent drug MTX in inhibiting Friend cells' growth. DP-MTX-71 was 4-fold more active compared to the α -MTX derivative, DP-MTX-93, which has the same bridging group.

Overall, under the conditions of the experimental system used, all tested DP-MTX derivatives, and especially DP-MTX-142, were less cytotoxic than the parent drug MTX (control). The tested alpha-DP-MTX derivatives were about 10- to 300-fold less toxic than MTX. These results indicate that alpha-DP-MTX derivatives are inactive prodrugs of the active species MTX.

Example 6: Effect of DP-MTX on DHFR activity

In order to further explore the cytotoxic effect of DP-MTX-71, the activity of this compound was measured on cell cultures of the human tumor cell CEM in the presence or absence of Leucovorin. Leucovorin is a reduced folate, used in chemotherapy with high doses of methotrexate for 'rescue' of host from toxicity.

Wild type human leukemia CEM cells or the MTX-resistant mutant cells, CEM-MTX, were plated into 96 well plates containing RPMI, 10% FCS at a concentration of 5×10^4 cells / ml. Either commercial MTX or the MTX derivative DP-MTX-71 was added at decreasing concentrations. Leucovorin, at a final concentration of 10 μ M, was added to some of the plates as indicated.

The plates were incubated for 72 hours (37°C, 5% CO₂) before being assayed for viability using the colorimetric MTT assay (Mosmann (1983) J. Immunol Methods 65: 55-63). EC₅₀ values (cytotoxic concentrations for 50% of cells in culture) were calculated for MTX and DP-MTX-71 in the presence and absence of Leucovorin.

Table 3: Effects of MTX and DP-MTX-71 on cell growth in the presence and absence of leucovorin

Compound	Cell line	EC ₅₀ , nMol No Leucovorin	EC ₅₀ , nMol +10 μ M Leucovorin
MTX	CEM	23 \pm 13	910 \pm 378
MTX	CEM-MTX	2535 \pm 224	19965 \pm 12807
DP-MTX-71	CEM	182 \pm 75	1817 \pm 639
DP-MTX-71	CEM-MTX	225 \pm 49	5798 \pm 3939

As can be seen from the results summarized in Table 3, the cytotoxic effect of DP-MTX-71 on the wild type and MTX-resistant human tumor cell lines was greatly reduced (10- and 26-folds, respectively) in the presence of the reduced folate Leucovorin. The fact that leucovorin presence interferes with the cytotoxic effect of

DP-MTX-71 is an indication that the drug inhibits the activity of dihydrofolate reductase (DHFR).

These results indicate that DP-MTX-71 inhibits DHFR, hence suggesting that it functions as a classical antifolate.

5

Example 7: Effect of DP-MTX-71 on PLA₂

DP-MTX-71 was evaluated for inhibition of phospholipase PLA₂-I and PLA₂-II activities (PanLab catalog # 160000 and 160100, respectively). Phospholipase PLA₂ activity was assayed in vitro as described by Katsumata et al. (Anal. Biochem. (1986) 154: 676-681) in the presence of 0.3, 3, 30 and 300 μ M DP-MTX-71. IC₅₀ values, namely the concentration of the tested compound that inhibits 50% of the enzyme activity, were calculated.

Greater than 50% inhibition was observed for the inhibition of PLA₂-I (IC₅₀ ~ 28 μ M). Moderate activity (41% inhibition at 300 μ M) was also noted for inhibition of PLA₂-II. These values are well within the range of IC₅₀ values calculated for other known inhibitors of PLA₂; NDGA and Quinacrine (IC₅₀ values 40 μ M and 390 μ M, respectively).

The fact that DP-MTX-71 inhibits PLA₂ indicates that the DP-MTX molecule may serve as a substrate for this enzyme.

20

Example 8: Effect of DP-MTX-71 on various tumor cell lines

The cytotoxic effect of DP-MTX-71 was evaluated on various tumor cell lines as indicated in Table 3 below. The cultured cells were incubated, during their linear growth phase, in the presence of increasing concentrations of either commercial MTX or the DP-MTX-71 derivative. Cell viability was analyzed after 72 hours incubation (37°C, 5% CO₂) using the colorimetric MTT assay (Mosmann (1983) J. Immunol Methods 65: 55-63). EC₅₀ values were calculated from the dose response curves and are summarized in Table 4.

30

Table 4: cytotoxic effect of DP-MTX-71 on various tumor cell lines

Name of cell line	MTX EC ₅₀ , nMol	DP-MTX-71 EC ₅₀ , nMol	Source
CEM	21	169	T-cell ALL (human)
C6	46	14	Glioma (rat)
Jurkat	21	162	T-cell lymphoma
Du148	29	16	Prostate cancer (human)
J774	112	22	Myeloma (mouse)
L1210	13	206	lymphocytic leukemia (mouse)
N87	13486	2332	Gastric carcinoma (human)
A431	850	737	Colon adenocarcinoma (human)
T47D	50000	23503	Mammary cancer (human)

As can be seen from the results presented in Table 4, the cell growth inhibitory effects of MTX and DP-MTX-71 greatly varied depending on the tested tumor cell line. MTX was found to be more effective than DP-MTX-71 in inhibiting cell growth of CEM, Jurkat and L1210 cell lines. On the other hand, DP-MTX-71 was more effective in inhibiting cell growth of C6 glioma, J774 myeloma and N87 carcinoma cell lines.

Example 9: DP-MTX-71 overcomes multi-drug resistance (MDR)

In order to further characterize the activity of DP-MTX-71, the cytotoxic effect of the compound was measured on cell cultures of drug resistant mutants derived from the mouse tumor cell line L1210.

Three cell lines were employed: the parent leukemia cell line which is sensitive to methotrexate (L1210), RFC-mediated MTX-resistant cell line (L1210-MTX), and a double mutated cell line which is resistant to MTX and Taxol (L1210-MTX-TAX). The Taxol resistance is MDR-mediated, due to increased activity of the P-glycoprotein (P-gp) pump. Cell death was tested using the MTT assay. EC50 values were calculated and the results are presented in Table 5.

Table 5: cytotoxic effect of DP-MTX-71 on mutant tumor cell lines

Cell line	MTX EC ₅₀ , nMol	DP-MTX-71 EC ₅₀ , nMol	TAX EC ₅₀ , nMol
L1210	14	207	8
L1210-MTX	1081	255	8
L1210-MTX-TAX	987	221	1846

5

As shown in Table 5, DP-MTX-71 was found to be more potent than the parent drug MTX in inhibiting cell growth of the MTX-resistant cell lines L1210-MTX, but not of the wild type cell L1210. This result is similar to the results obtained with the human leukemia cell line CEM and the MTX-resistant cell line CEM-MTX.

10 In addition, DP-MTX-71 was found to be about 4-5 times more potent than MTX in inhibiting growth of the double mutant L1210-MTX-TAX that is characterized by increased activity of P-gp.

These results indicate that DP-MTX-71 may overcome multi-drug resistance (MDR).

15 **Example 10: DP-MTX-71 does not provoke MDR phenotype**

An effort was directed towards generation of DP-MTX-71 resistant cell line. The following protocol for mutant selection was followed. Exponentially growing Chinese hamster ovary (CHO) AA8 cells in monolayer conditions were washed with PBS, trypsinized and plated (300,000 cells/T25 flask) in alpha MEM medium (Beth Haemek, Israel) containing 5% fetal calf serum, penicillin/streptomycin and 2 mM glutamine. DP-MTX-71 (from a 1 mM stock) was added at a 50 nM concentration at the time of cell seeding. Ten to fourteen days later, when cells doubled their density, the attached cells were washed with PBS, trypsinized, viability counted, and plated as above in the presence of 50 nM and 75 nM DP-MTX-71 (50% increment).

20

This multiple step drug selection was continued by using 25-50% increments in drug concentration over a period of 8 months. Selection was terminated at 1 micromolar DP-MTX-71 concentration.

Only a single resistant cell line was obtained after eight months of repeated
5 passage of the cells under the selective pressure of presence of DP-MTX-71. This period is significantly longer than the time required to obtain cell lines resistant to the parent drug MTX or other known derivatives thereof. The obtained DP-MTX-71 resistant cell line was characterized and was found to acquire resistance by over-expressing the target enzyme DHFR. It is important to note that DP-MTX-71 did not
10 induce over-expression of P-gp. This may be an indication that P-gp does not affect DP-MTX-71.

From these results it appears that DP-MTX-71 has a potential to serve as a cytotoxic agent that lacks or has a reduced ability to provoke multi-drug resistance (MDR).

15 **Example 11: Effect of DP-MTX on survival of tumor bearing mice (viability studies)**

The anti-tumor effects of DP-MTX were evaluated in vivo on mice transplanted with L1210 murine leukemia cells.

Mice of strain B6D2F1/J (stock number 100006), which are F1 cross of
20 C57/BJ x DBA/2J, were purchased from The Jackson Laboratory (JAX Mice & Mouse Model Related Services, NH, USA). Animals were allowed 10 days adjustment time at our animal care facility before initiation of the experiment.

Mice were transplanted with 1×10^6 L1210 (mouse leukemia; ATCC # CCL-219) or L1210/MTX (RFC-mediated MTX resistant L1210 mutant; provided by Dr.
25 Y. Assraf, Technion, Israel) cells by injection into the peritoneum (i.p.) (day 1). Two days later (on day 3) groups of animals (8 mice per group) were injected i.p. with either 3 mg/Kg commercial MTX (Abitrexate®, from ABIC, Israel) or molar equivalent dose of DP-MTX-71. Another group of 8 animals were injected with vehicle solution (2% propylene-glycol, 0.2% Tween-80 in PBS) and serve as control
30 group. Thereafter, the same doses of MTX or DP-MTX-71 (MTX equivalent dose) or

vehicle only, were injected i.p. every two days until day 17. Mice survival was monitored daily until the end of the experiment on day 21.

As shown in Figure 1, mice implanted i.p. with tumors died after about 10 days. MTX treatment, which by itself caused no mortality in the mice, effectively prolonged the life of mice implanted with tumors; in this case animal death started on day 15. Treatment with DP-MTX-71 was the most effective, and prolonged survival of animals transplanted with L1210 tumor cells for at least 7 days beyond survival of animals treated with MTX. At the end of the experiment, on day 21, only one mouse (out of 8 mice) survived in the control group or in the group treated with MTX, in comparison to five mice (out of 8) survived in the group treated with DP-MTX-71.

Similar trend (results not shown) was observed in the experiment where the mice were transplanted with L1210/MTX cells (tumor cell line resistant to MTX). In this case treatment with 3 mg/Kg dose equivalent DP-MTX-71 prolonged animal survival by 3 days beyond survival of animals treated with MTX. Furthermore, a dose dependent effect of DP-MTX-71 was observed. Prolongation of animal survival for 17 days beyond survival of animals treated with MTX was observed for 50% of the mice treated with 10 mg/Kg dose equivalent of DP-MTX-71 (i.p. injected every 4 days).

These results indicate that DP-MTX-71 is more effective than the patent drug MTX in increasing life span of tumor bearing mice.

Example 12: Effect of DP-MTX in preventing tumor growth (in vivo)

Protection against tumor growth was examined in mice pretreated with MTX or DP-MTX-71.

Female B6D2F1/J mice were injected i.p. with either 3 mg/Kg commercial MTX (6 animals in a group) or equivalent dose of DP-MTX-71 (8 animals in a group) on every other day for a period of 21 days. At the end of this regimen, the mice were injected with 1×10^6 L1210 cells into the peritoneum. The animals were then monitored for viability for 12 days subsequent to the injection of the tumor cells. As can be seen in Table 6, at the end of the experiment, namely 12 days after the

injection of the tumor cells, 2 and 5 animals survived in the groups treated, respectively, with MTX and DP-MTX-71.

5 **Table 6: Effect of DP-MTX on survival of mice implanted with L1210 tumor cells**

Days after tumor implantation	Surviving mice (MTX pre-treatment)	Surviving mice (DP-MTX-71 pre-treatment)
9	6/6	8/8
10	4/6	8/8
11	3/6	8/8
12	2/6	5/8

10 DP-MTX derivative shows an improved pharmacokinetic properties comparing to the parent drug MTX. Pre-treatment with DP-MTX-71 was effective in increasing survival of mice implanted with tumor cells.

15 **Example 13: Anti-inflammatory activity of DP-MTX (in the carrageenan-induced rat paw oedema model)**

 The potential anti-inflammatory activities of various DP-MTX prodrugs are evaluated by employing the experimental model of carrageenan-induced paw oedema in the rat.

20 Carrageenan-induced rat paw edema is a widely employed animal model for acute inflammation. The objective of the study is to assess the potential prophylactic effects of DP-MTX derivatives on the prevention of inflammatory swelling and, in particular, to compare the efficacy parameters with those obtained for MTX. The experimental set-up is as follows: Male Sprague-Dawley rats weighing 120-180g
25 (supplied by Harlan Laboratories Breeding Center, Israel) are intraperitoneally (i.p.)

injected with MTX or DP-MTX one hour prior to the induction of inflammation with carrageenan. MTX and the tested MTX-derivatives are applied at a dose level of 1 mg/kg body weight. Animals injected with the vehicle only serve as a control group.

Paw edema is induced by a single sub-plantar injection of 0.1 ml 2% carrageenan in physiological saline, into the right hind paw of each test animal. Just prior to paw edema induction, the paw thickness of the test animals' right hind paws is measured in duplicate using a plethysmograph and micrometer to provide a baseline. At 3, 5 and 7 hours post carrageenan injection, the right paw thickness is measured in the same manner as before.

The assessment of potential anti-inflammatory activity is based on the relative reduction (%) in mean group values of carrageenan-induced right hind paw edema in animal groups treated with tested compounds versus respective values in the control group treated with vehicle only. The effect of DP-MTX on carrageenan-induced paw edema is evaluated in comparison to the effect of methotrexate by following severity of inflammation as determined by hind paw swelling.

Studies of this kind indicate that the phospholipid derivatives of methotrexate have higher therapeutic indices compared to methotrexate in the carrageenan-induced paw edema model for inflammation.

Example 14: Effect of DP-MTX-71 in collagen arthritis inflammation model

The potential effect of DP-MTX-71 in treatment of arthritis was examined in collagen arthritis model of inflammation in mice.

Groups of 5 Balb/cByJ mice strain, 6-8 weeks of age, were used for the induction of arthritis by monoclonal antibodies (mAbs) plus lipopolysaccharide (LPS). The animals were administered intravenously a combination of 4 different mAbs (PanLab's protocol # 553600) in total of 4 mg/mouse (day 0), followed by intravenous 25 µg of LPS 72 hours later (day 3). From day 3, one hour after LPS administration, DP-MTX-71 (30 mg/kg) was administered orally once daily for 3 consecutive days. Animals treated with vehicle serve as control group. Paw edema was monitored by using a plethysmometer for measuring paw volume of the two hind paws on day 0, 5, 7, 10, 14 and 17. Assessment of anti-inflammatory activity was

based on the relative reduction (%) in hind paw swelling in animals treated with tested compound in comparison to the control group treated with vehicle only.

Results: significant reduction of hind paws swelling of 22%, 27%, 27% and 30% was noted on days 5, 7, 10 and 17, respectively.

- 5 DP-MTX-71 shows anti-inflammatory activity as demonstrated in the collagen arthritis animal model.

Example 15: *In vivo* efficacy and toxicity studies of DP-MTX (in rat adjuvant-induced arthritis)

- 10 DP-MTX compounds were evaluated for efficacy and toxicological profile in the progression of adjuvant-induced arthritis (AA) in Lewis rats, which is a widely employed animal model for the human rheumatoid arthritis (RA). The effects of DP-MTX were compared with those obtained with MTX.

- 15 Arthritis was induced in the left hind paws of male Lewis rats by sub-plantar injection of a suspension of *Mycobacterium butyricum* in paraffin oil. Test animals were intraperitoneally (i.p.) injected with either 0.02 mg/kg or 0.1 mg/kg of DP-MTX prodrug or the commercially available methotrexate (Abitrexate®). Dosing was initiated on the day of AA induction and thereupon continued as once-a-day repeated injections throughout a 3-week study period. An additional group of rats was treated
20 with solvent only (1% propylene glycol and 0.01% Tween-80 in USP water) and served as a control group.

- Assessment of the DP-MTX effects was based on relative changes in hind paw thickness, measured prior to the treatment, during and at termination of the study period. These effects were compared to the respective effects of MTX. Determination
25 of body weight gain at study termination was used as a measure of general systemic anti-inflammatory activity. In addition, all tested animals were inspected for clinical signs (e.g. histopathological patterns in the spleen, liver and intestines) to evaluate toxicity.

- Results indicate that phospholipid derivatives of methotrexate are effective in
30 suppression of AA-induced inflammatory phenotypes such as swelling, and evoke less pathological events.

Example 16: In vivo efficacy studies of DP-MTX (in the experimental allergic encephalomyelitis (EAE) model system)

The effects of DP-MTX compounds on sites of inflammation within the central nervous system are assessed in an animal model. The chosen model system is
5 experimental allergic encephalomyelitis (EAE) which is a known animal model for multiple sclerosis (MS).

In a preliminary set of experiments EAE was induced in SJL/J mice by either one of two schemes: a) active induction by injecting mice via the footpads with proteolipid protein (PLP₁₃₉₋₁₅₁) emulsified in complete Freund's adjuvant (CFA) as
10 described by Weinberg et al. (J. Immunol. (1999) 162: 1818-1826), or b) passive induction, namely adoptive transfer of EAE accomplished by transferring reactive cells from a donor. Briefly, PLP-reactive lymph node cells are drawn from mice inoculated with PLP-CFA, 10 days post inoculation. The cells are incubated in vitro for 4 days with soluble PLP, then collected, washed and transferred to naive mice.

15 The tested DP-MTX compounds are injected either i.p or i.v. at 8, 10 or 12 days post inoculation in the case of the active induction of EAE, or at days 2, 4 and 6 after cell transfer in the passive induction protocol. According to both procedures, the animals are sacrificed 4 hours after injection of the tested DP-MTX compound. Blood samples are collected and the animals' brains and spinal cords are excised and
20 homogenized. The levels of MTX in serum and tissue extracts of the treated animals are determined. Briefly, samples of brain or spinal cord are suspended in three volumes of 5 mg/ml sodium ascorbate. Samples are homogenized, sonicated and centrifuged at 27000g for 30 min. Supernatants are placed in boiling water bath for 10 min. and then centrifuged at 3000g for 1 min. The obtained clear supernatants are
25 used for determination of free MTX. Abbot's TDX method and kit are used to determine the levels of MTX in serum and tissue extracts.

A group of mice with EAE are kept for a longer time while receiving leucovorin daily and is watched for EAE symptoms.

EAE severity is evaluated statistically by following two main parameters:

a) neurological symptoms scored according to a standard score in which: 0- no clinical signs; 1- limp tail; 2- ataxia; 3- hind limb paralysis; 4- complete paralysis and incontinence; 5- moribund and death; and

5 b) histopathological symptoms which are evaluated by using sections of the brain stained with hematoxylin-eosin for detection of mononuclear cell infiltration and with luxol-fast for the detection of demyelination.

Results have shown that the tested DP-MTX compounds reach the site of inflammation and exert their therapeutic effect in inhibiting the clinical signs of EAE. Thus, DP-MTX may be useful in treatment of CNS autoimmune diseases and
10 inflammation.

Example 17: Effects of DP-MTX and DP-5FUdR compounds in restenosis model system in rats.

The anti-proliferative effects of compounds of the invention are assessed in a
15 model system for restenosis in rats. Restenosis is a phenomenon of neointimal formation occurring following percutaneous transluminal coronary angioplasty (PTCA) procedures.

Experimental neointimal formation is induced by balloon injury in rats according to the following procedure. Male Sprague-Dawley rats (380-450 g) are
20 anesthetized by inhalation of halothane and dinitric oxyde. After the right common carotid artery and the right external carotid artery are exposed, a 2F Fogarty arterial embolectomy catheter (Baxter Healthcare, Santa Ana, CA) is inserted into the lumen of the right external carotid artery and is guided to a fixed distance (about 5 cm). The balloon is inflated with saline and is withdrawn at a constant rate back to a point
25 proximal to the site of insertion. This procedure is repeated three times.

The effect of the balloon injury is assessed 2 to 4 weeks following the above-described treatment. The left (control, untreated) and right (experiment, treated) carotid arteries of rats under penthal overdose, are perfused in situ with 4% formaline, pH 7.4 and then excised. Each of the left and right arteries is cut at three segments:
30 upper, middle and lower, and then fixed in formaline. Cross-sections are stained with van Gieson's stain and the intima and media thickness is evaluated under a

microscope. A higher calculated ratio of intima/media in the treated (right) artery in comparison to the control (left) artery, is an indication of neointimal formation and development of restenosis.

Various DP-MTX and DP-5-FUdR compounds are tested for preventive and/or
5 curative effects on restenosis by carrying-out the following schemes:

- a) Preventive procedure – the tested compound is daily administered starting from 7 to 3 days prior to the induced balloon injury, and up to the day of the induced injury.
- b) Curative procedure – the tested compound is daily administered starting from
10 the day of induced balloon injury and up to 14 to 30 days after the induced injury.

Peroral- and parenteral- , such as subcutaneous, modes of administration may be employed in both the preventive and curative procedures.

In yet another regimen, a combination of the preventive and the curative procedures is
15 tested in the rat model system for restenosis.


Both DP-MTX and DP-5-FUdR compounds are promising agents for inhibition of intimal proliferation and may be useful in clinical procedures.

20 While the present invention has been particularly described, persons skilled in the art will appreciate that many variations and modifications can be made. Therefore, the invention is not to be construed as restricted to the particularly described embodiments, rather the scope, spirit and concept of the invention will be more readily understood by reference to the claims which follow.

25

CLAIMS:

1. A prodrug of the general formula I:



5

Formula I

or a pharmaceutically acceptable salt thereof, wherein:

R₁ is a saturated or unsaturated, straight-chain or branched, substituted or unsubstituted hydrocarbon chain having from 2 to 30 carbon atoms;

- 10 **R₂** is H or a phospholipid head group;

Z is saturated or unsaturated, straight-chain or branched, substituted or unsubstituted hydrocarbon chain having from 2 to 15 carbon atoms, which may include cyclic elements, and optionally is interrupted by one or more atoms selected from oxygen and sulfur atoms;

- 15 **X** is a direct covalent bond or selected from the group consisting of O, S, NH and C(O) groups; and

D is a residue of an anti-proliferative drug,

wherein the bound anti-proliferative drug residue is an inactive form of the drug which is selectively activated in cells and tissues with elevated phospholipase activity.

20

2. The prodrug according to claim 1, wherein the anti-proliferative drug is methotrexate or pharmaceutically acceptable derivatives thereof.

3. The prodrug according to claim 1, wherein the anti-proliferative drug is 2'-deoxy-5-fluorouridine and pharmaceutically acceptable derivatives thereof.

4. The prodrug according to claim 1, wherein an ester bond at position
5 sn-2 of the phospholipid of the general formula I is cleaveable by a phospholipase.

5. The prodrug according to claim 4, wherein said phospholipase is phospholipase A₂ (PLA₂).

10 6. The prodrug according to claim 1, wherein R1 is a hydrocarbon chain having from 5 to 20 carbon atoms.

7. The prodrug according to claim 1, wherein R1 is a hydrocarbon chain having 15 or 17 carbon atoms.

15

8. The prodrug according to claim 1, wherein R2 is selected from the group consisting of choline, ethanolamine, inositol and serine.

9. The prodrug according to claim 1 selected from the group consisting of:

20 1-Stearoyl-2-[3-(α -MTX amido)-Propanoyl]-sn-Glycero-3-Phosphatidylcholine,

1-Stearoyl-2-[3-(γ -dodecylate- α -MTX amido)-Propanoyl]-sn-Glycero-3-Phosphatidylcholine,

25 1-Stearoyl-2-[4-(α -MTX amido)-Butanoyl]-sn-Glycero-3-Phosphatidylcholine,

1-Stearoyl-2-[6-(α -MTX amido)-Hexanoyl]-sn-Glycero-3-Phosphatidylcholine,

1-Stearoyl-2-[8-(α -MTX amido)-Octanoyl]-sn-Glycero-3-Phosphatidylcholine,

30 1-Stearoyl-2-[8-(γ -dodecylate- α -MTX amido)-Octanoyl]-sn-Glycero-3-Phosphatidylcholine,

1-Stearoyl-2-[3-(α -dodecylate- γ -MTX amido)-Propanoyl]-sn-Glycero-3-Phosphatidylcholine, and

1-stearoyl-2-[5^m-(2''-deoxy-5'-fluorouridine-5''-)-3^m,3^m-dimethyl] glutaroyl-1^m-sn-glycero-3-phosphatidylcholine

5

10. The prodrug according to claim 1 which is 1-Stearoyl-2-[3-(α -MTX amido)-Propanoyl]-sn-Glycero-3-Phosphatidylcholine.

11. The prodrug according to claim 1 which is 1-Stearoyl-2-[3-(α -dodecylate- γ -MTX amido)-Propanoyl]-sn-Glycero-3-Phosphatidylcholine.

10

12. A pharmaceutical composition comprising, as an active ingredient, a prodrug of the general formula I according to any one of claims 1 to 11, and a pharmaceutically acceptable carrier.

15

13. The pharmaceutical composition according to claim 12 further comprising an additional anti-neoplastic agent.

14. The pharmaceutical composition according to claim 12 or claim 13, which is suitable for oral, ocular, nasal, parenteral, topical or rectal administration.

20

15. The pharmaceutical composition according to claim 12 or claim 13, which is suitable for oral administration, intravenous administration or topical administration.

25

16. The pharmaceutical composition according to claim 12 or claim 13, in the form of solutions, suspensions, capsules, tablets, aerosols, gels, ointments or suppositories.

17. Use of a prodrug as defined in any one of claims 1 to 11 for the manufacture of a medicament.

30

18. A method for treatment of a disease or disorder related to an inflammatory condition comprising administering to a patient in need thereof a therapeutically effective amount of a pharmaceutical composition according to claim 12.

19. The method according to claim 18, wherein said disease or disorder related to an inflammatory condition is selected from the group consisting of granulomatous diseases, arthritis, rheumatoid arthritis, multiple sclerosis, systemic sclerosis, asthma, psoriasis, systemic lupus erythematosus, inflammatory bowel syndromes and migraines.

20. A method for treatment of a disease or disorder related to a degenerative or atrophic condition comprising administering to a patient in need thereof a therapeutically effective amount of a pharmaceutical composition according to claim 12.

21. The method according to claim 20, wherein said disease or disorder related to a degenerative or atrophic condition is a central or peripheral neurological disease or disorder.

22. The method according to claim 20, wherein said disease or disorder related to a degenerative or atrophic condition is selected from the group consisting of autoimmune, cerebrovascular and neurodegenerative diseases and disorders such as idiopathic dementia, vascular dementia, multi-infarct dementia, traumatic dementia, Alzheimer's disease, Pick's disease, Huntington's disease, dementia paralitica, Parkinson's disease, diabetic neuropathy, amyotrophic lateral sclerosis, ischemia of the optic nerve, age-related macular degeneration, stroke and trauma.

23. A method for treatment of a disease or disorder related to uncontrolled cell growth comprising administering to a patient in need thereof a therapeutically effective amount of a pharmaceutical composition according to claim 12.

5 24. The method according to claim 23, wherein said disease or disorder related to uncontrolled cell growth is a neoplastic growth.

25. The method according to claim 24 wherein said neoplastic growth is a primary or a secondary tumor.

10 26. The method according to claim 24 wherein said neoplastic growth is a drug-resistant tumor.

15 27. The method according to claim 24 wherein said neoplastic growth is a methotrexate-resistant tumor.

28. The method according to claim 24 wherein said neoplastic growth is a multidrug-resistant tumor.

20 29. The method according to claim 23, wherein said disease or disorder related to uncontrolled cell growth is selected from the group consisting of psoriasis, lymphocytic leukemia, myelogenous leukemia, Burkitt's lymphoma, non-Hodgkin's lymphomas, mycosis fungoides, osteosarcoma, hydatidiform mole, trophoblastic diseases such as chorioadenoma destruens and choriocarcinoma, and carcinomas of
25 the head and neck, breast, liver, lung, colon, ovary, cervix, urinary, bladder, prostate, pancreas, skin, the gastrointestinal tract and the oropharyngeal areas.

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
22 March 2001 (22.03.2001)

PCT

(10) International Publication Number
WO 01/19320 A3

(51) International Patent Classification⁷: **A61K 31/56**

(21) International Application Number: **PCT/IL00/00562**

(22) International Filing Date:
13 September 2000 (13.09.2000)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
131887 14 September 1999 (14.09.1999) IL

(71) Applicant (for all designated States except US):
D-PHARM LIMITED [IL/IL]; Kiryat Weizmann
Science Park, P.O. Box 2313, 76123 Rehovot (IL).

(72) Inventors; and

(75) Inventors/Applicants (for US only): **KOZAK, Alexan-**
der [IL/IL]; 8 Naphtali Ben Ephraim Street, 76555 Re-
hovot (IL). **SHAPIRO, Israel** [IL/IL]; 1 Aharon Bouga-
nium Street, 75122 Ramla (IL). **VINNIKOVA, Marina**

[IL/IL]; 1 Aharon Bouganium Street, 75122 Ramla (IL).
ERSHOV, Leonid [IL/IL]; Ganei Shemesh 64/20, 99510
Beit Shemesh (IL). **SENDERIKHIN, Alexander** [IL/IL];
Eben Ezra 11/32, 77622 Ashdod (IL). **AYALON, Oran**
[IL/IL]; Shikun Banim 143, 76922 Ganei Yohanan (IL).


(74) Agent: **WALKER, Naomi**; Kiryat Weizmann Science
Park, P.O. Box 2313, 76123 Rehovot (IL).

(81) Designated States (national): AE, AG, AL, AM, AT, AU,
AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CR, CU, CZ,
DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR,
HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR,
LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ,
NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM,
TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.

(84) Designated States (regional): ARIPO patent (GH, GM,
KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian
patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European
patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE,
IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG,
CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

[Continued on next page]

(54) Title: PHOSPHOLIPID PRODRUGS OF ANTI-PROLIFERATIVE DRUGS



(57) Abstract: The invention discloses prodrugs comprising anti-proliferative drugs covalently linked, via a bridging group, to a phospholipid moiety such that the active species is preferentially released, preferably by enzymatic cleavage, at the required site of action. The invention further discloses pharmaceutical compositions said prodrugs and the uses thereof for the treatment of diseases and disorders related to inflammatory, to degenerative or atrophic conditions, and to uncontrolled cell growth. Fig. 1 depicts a graph of animal survival during the course of an experiment wherein mice were i.p. transplanted with 11210 mouse leukemia cells and then treated with vehicle only (squares), MTX (triangles) or molar equivalent dose of DP-MTX071 (circles) according to the regimen described in example in Example 11.



Published:

— with international search report

(88) Date of publication of the international search report:
27 September 2001

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

PCT

For r

Office use only 10/088160

REQUEST

The undersigned requests that the present international application be processed according to the Patent Cooperation Treaty.

International Application No.

International Filing Date

Name of receiving Office and "PCT International Application"

Applicant's or agent's file reference
(if desired) (12 characters maximum) DPL/009/PCT

Box No. I TITLE OF INVENTION

PHOSPHOLIPID PRODRUGS OF ANTI-PROLIFERATIVE DRUGS

Box No. II APPLICANT

Name and address: (Family name followed by given name; for a legal entity, full official designation. The address must include postal code and name of country. The country of the address indicated in this Box is the applicant's State (that is, country) of residence if no State of residence is indicated below.)

D-PHARM LIMITED
KIRYAT WEIZMANN SCIENCE PARK
REHOVOT 76326
ISRAEL

☐ This person is also inventor.

Telephone No.

972-8-9300794

Facsimile No.

972-8-9300794

Teleprinter No.

State (that is, country) of nationality:

IL

State (that is, country) of residence:

IL

This person is applicant
for the purposes of:☐all designated
States☒all designated States except
the United States of America☐the United States
of America only☐the States indicated in
the Supplemental Box

Box No. III FURTHER APPLICANT(S) AND/OR (FURTHER) INVENTOR(S)

Name and address: (Family name followed by given name; for a legal entity, full official designation. The address must include postal code and name of country. The country of the address indicated in this Box is the applicant's State (that is, country) of residence if no State of residence is indicated below.)

KOZAK, ALEXANDER,
8 NAPHTALI BEN EPHRAIM ST.
REHOVOT 76555
ISRAEL

This person is:

☐ applicant only☒ applicant and inventor☐ inventor only (If this check-box
is marked, do not fill in below.)

State (that is, country) of nationality:

IL

State (that is, country) of residence:

IL

This person is applicant
for the purposes of:☐all designated
States☐all designated States except
the United States of America☒the United States
of America only☐the States indicated in
the Supplemental Box☒ Further applicants and/or (further) inventors are indicated on a continuation sheet.

Box No. IV AGENT OR COMMON REPRESENTATIVE; OR ADDRESS FOR CORRESPONDENCE

The person identified below is hereby/has been appointed to act on behalf of the applicant(s) before the competent International Authorities as:

☒

agent

☐

common representative

Name and address: (Family name followed by given name; for a legal entity, full official designation. The address must include postal code and name of country.)

WALKER, NAOMI
KIRYAT WEIZMANN SCIENCE PARK
P.O. BOX 2313
REHOVOT 76123
ISRAEL

Telephone No.

972-8-9300794

Facsimile No.

972-8-9300795

Teleprinter No.

☐ Address for correspondence: Mark this check-box where no agent or common representative is/has been appointed and the space above is used instead to indicate a special address to which correspondence should be sent.

Continuation of Box No. III FURTHER APPLICANT(S) AND/OR (FURTHER) INVENTOR(S)

If none of the following sub-boxes is used, this sheet should not be included in the request

Name and address: (Family name followed by given name; for a legal entity, full official designation. The address must include postal code and name of country. The country of the address indicated in this Box is the applicant's State (that is, country) of residence if no State of residence is indicated below.)

SHAPIRO, ISRAEL
1 AHARON BOUGANIUM St.
RAMLA 75122
ISRAEL

This person is:

- ☐ applicant only
☒ applicant and inventor
☐ inventor only (If this check-box is marked, do not fill in below.)

State (that is, country) of nationality:
IL

State (that is, country) of residence:
IL

This person is applicant for the purposes of:

- ☐ all designated States ☐ all designated States except the United States of America ☒ the United States of America only ☐ the States indicated in the Supplemental Box

Name and address: (Family name followed by given name; for a legal entity, full official designation. The address must include postal code and name of country. The country of the address indicated in this Box is the applicant's State (that is, country) of residence if no State of residence is indicated below.)

VINNIKOVA, MARINA
1 AHARON BOUGANIUM St.
RAMLA 75122
ISRAEL

This person is:

- ☐ applicant only
☒ applicant and inventor
☐ inventor only (If this check-box is marked, do not fill in below.)

State (that is, country) of nationality:
IL

State (that is, country) of residence:
IL

This person is applicant for the purposes of:

- ☐ all designated States ☐ all designated States except the United States of America ☒ the United States of America only ☐ the States indicated in the Supplemental Box

Name and address: (Family name followed by given name; for a legal entity, full official designation. The address must include postal code and name of country. The country of the address indicated in this Box is the applicant's State (that is, country) of residence if no State of residence is indicated below.)

ERSHOV, LEONID
GANEI SHEMESH 64/20
BEIT SHEMESH 99510
ISRAEL

This person is:

- ☐ applicant only
☒ applicant and inventor
☐ inventor only (If this check-box is marked, do not fill in below.)

State (that is, country) of nationality:
IL

State (that is, country) of residence:
IL

This person is applicant for the purposes of:

- ☐ all designated States ☐ all designated States except the United States of America ☒ the United States of America only ☐ the States indicated in the Supplemental Box

Name and address: (Family name followed by given name; for a legal entity, full official designation. The address must include postal code and name of country. The country of the address indicated in this Box is the applicant's State (that is, country) of residence if no State of residence is indicated below.)

SENDERICHIN, ALEXANDER
EBEN EZRA 11/32
ASHDOD 77622
ISRAEL

This person is:

- ☐ applicant only
☒ applicant and inventor
☐ inventor only (If this check-box is marked, do not fill in below.)

State (that is, country) of nationality:
IL

State (that is, country) of residence:
IL

This person is applicant for the purposes of:

- ☐ all designated States ☐ all designated States except the United States of America ☒ the United States of America only ☐ the States indicated in the Supplemental Box

☐ Further applicants and/or (further) inventors are indicated on another continuation sheet.

Box No.V DESIGNATING STATES

The following designations are hereby made under Rule 4.9(a) (mark the applicable check-boxes, at least one must be marked):

Regional Patent

- ☒ **AP ARIPO Patent:** GH Ghana, GM Gambia, KE Kenya, LS Lesotho, MW Malawi, MZ Mozambique, SD Sudan, SL Sierra Leone, SZ Swaziland, TZ United Republic of Tanzania, UG Uganda, ZW Zimbabwe, and any other State which is a Contracting State of the Harare Protocol and of the PCT
- ☒ **EA Eurasian Patent:** AM Armenia, AZ Azerbaijan, BY Belarus, KG Kyrgyzstan, KZ Kazakhstan, MD Republic of Moldova, RU Russian Federation, TJ Tajikistan, TM Turkmenistan, and any other State which is a Contracting State of the Eurasian Patent Convention and of the PCT
- ☒ **EP European Patent:** AT Austria, BE Belgium, CH and LI Switzerland and Liechtenstein, CY Cyprus, DE Germany, DK Denmark, ES Spain, FI Finland, FR France, GB United Kingdom, GR Greece, IE Ireland, IT Italy, LU Luxembourg, MC Monaco, NL Netherlands, PT Portugal, SE Sweden, and any other State which is a Contracting State of the European Patent Convention and of the PCT
- ☒ **OA OAPI Patent:** BF Burkina Faso, BJ Benin, CF Central African Republic, CG Congo, CI Côte d'Ivoire, CM Cameroon, GA Gabon, GN Guinea, GW Guinea-Bissau, ML Mali, MR Mauritania, NE Niger, SN Senegal, TD Chad, TG Togo, and any other State which is a member State of OAPI and a Contracting State of the PCT (if other kind of protection or treatment desired, specify on dotted line)


National Patent (if other kind of protection or treatment desired, specify on dotted line):

- | | |
|--|--|
| <input checked="" type="checkbox"/> AE United Arab Emirates | <input checked="" type="checkbox"/> LC Saint Lucia |
| <input checked="" type="checkbox"/> AG Antigua and Barbuda | <input checked="" type="checkbox"/> LK Sri Lanka |
| <input checked="" type="checkbox"/> AL Albania | <input checked="" type="checkbox"/> LR Liberia |
| <input checked="" type="checkbox"/> AM Armenia | <input checked="" type="checkbox"/> LS Lesotho |
| <input checked="" type="checkbox"/> AT Austria | <input checked="" type="checkbox"/> LT Lithuania |
| <input checked="" type="checkbox"/> AU Australia | <input checked="" type="checkbox"/> LU Luxembourg |
| <input checked="" type="checkbox"/> AZ Azerbaijan | <input checked="" type="checkbox"/> LV Latvia |
| <input checked="" type="checkbox"/> BA Bosnia and Herzegovina | <input checked="" type="checkbox"/> MA Morocco |
| <input checked="" type="checkbox"/> BB Barbados | <input checked="" type="checkbox"/> MD Republic of Moldova |
| <input checked="" type="checkbox"/> BG Bulgaria | <input checked="" type="checkbox"/> MG Madagascar |
| <input checked="" type="checkbox"/> BR Brazil | <input checked="" type="checkbox"/> MK The former Yugoslav Republic of Macedonia |
| <input checked="" type="checkbox"/> BY Belarus | <input checked="" type="checkbox"/> MN Mongolia |
| <input checked="" type="checkbox"/> BZ Belize | <input checked="" type="checkbox"/> MW Malawi |
| <input checked="" type="checkbox"/> CA Canada | <input checked="" type="checkbox"/> MX Mexico |
| <input checked="" type="checkbox"/> CH and LI Switzerland and Liechtenstein | <input checked="" type="checkbox"/> MZ Mozambique |
| <input checked="" type="checkbox"/> CN China | <input checked="" type="checkbox"/> NO Norway |
| <input checked="" type="checkbox"/> CR Costa Rica | <input checked="" type="checkbox"/> NZ New Zealand |
| <input checked="" type="checkbox"/> CU Cuba | <input checked="" type="checkbox"/> PL Poland |
| <input checked="" type="checkbox"/> CZ Czech Republic | <input checked="" type="checkbox"/> PT Portugal |
| <input checked="" type="checkbox"/> DE Germany | <input checked="" type="checkbox"/> RO Romania |
| <input checked="" type="checkbox"/> DK Denmark | <input checked="" type="checkbox"/> RU Russian Federation |
| <input checked="" type="checkbox"/> DM Dominica | <input checked="" type="checkbox"/> SD Sudan |
| <input checked="" type="checkbox"/> DZ Algeria | <input checked="" type="checkbox"/> SE Sweden |
| <input checked="" type="checkbox"/> EE Estonia | <input checked="" type="checkbox"/> SG Singapore |
| <input checked="" type="checkbox"/> ES Spain | <input checked="" type="checkbox"/> SI Slovenia |
| <input checked="" type="checkbox"/> FI Finland | <input checked="" type="checkbox"/> SK Slovakia |
| <input checked="" type="checkbox"/> GB United Kingdom | <input checked="" type="checkbox"/> SL Sierra Leone |
| <input checked="" type="checkbox"/> GD Grenada | <input checked="" type="checkbox"/> TJ Tajikistan |
| <input checked="" type="checkbox"/> GE Georgia | <input checked="" type="checkbox"/> TM Turkmenistan |
| <input checked="" type="checkbox"/> GH Ghana | <input checked="" type="checkbox"/> TR Turkey |
| <input checked="" type="checkbox"/> GM Gambia | <input checked="" type="checkbox"/> TT Trinidad and Tobago |
| <input checked="" type="checkbox"/> HR Croatia | <input checked="" type="checkbox"/> TZ United Republic of Tanzania |
| <input checked="" type="checkbox"/> HU Hungary | <input checked="" type="checkbox"/> UA Ukraine |
| <input checked="" type="checkbox"/> ID Indonesia | <input checked="" type="checkbox"/> UG Uganda |
| <input checked="" type="checkbox"/> IL Israel | <input checked="" type="checkbox"/> US United States of America |
| <input checked="" type="checkbox"/> IN India | <input checked="" type="checkbox"/> UZ Uzbekistan |
| <input checked="" type="checkbox"/> IS Iceland | <input checked="" type="checkbox"/> VN Viet Nam |
| <input checked="" type="checkbox"/> JP Japan | <input checked="" type="checkbox"/> YU Yugoslavia |
| <input checked="" type="checkbox"/> KE Kenya | <input checked="" type="checkbox"/> ZA South Africa |
| <input checked="" type="checkbox"/> KG Kyrgyzstan | <input checked="" type="checkbox"/> ZW Zimbabwe |
| <input checked="" type="checkbox"/> KP Democratic People's Republic of Korea | |
| <input checked="" type="checkbox"/> KR Republic of Korea | |
| <input checked="" type="checkbox"/> KZ Kazakhstan | |

Check-box reserved for designating States which have become party to the PCT after issuance of this sheet:

☒

Precautionary Designation Statement: In addition to the designations made above, the applicant also makes under Rule 4.9(b) all other designations which would be permitted under the PCT except any designation(s) indicated in the Supplemental Box as being excluded from the scope of this statement. The applicant declares that those additional designations are subject to confirmation and that any designation which is not confirmed before the expiration of 15 months from the priority date is to be regarded as withdrawn by the applicant at the expiration of that time limit. (Confirmation (including fees) must reach the receiving Office within the 15-month time limit.)

Box No. VI PRIORITY CLAIM		<input type="checkbox"/> Further priority claim indicated in the Supplemental Box.		
Filing date of earlier application (day/month/year)	Number of earlier application	Where earlier application is:		
		national application: country	regional application: regional Office	international application: receiving Office
item (1) 14 September 1999 (14.09.1999)	131887	IL		
item (2)				
item (3)				
<input checked="" type="checkbox"/> The receiving Office is requested to prepare and transmit to the International Bureau a certified copy of the earlier application(s) (only if the earlier application was filed with the Office which for the purposes of the present international application is the receiving Office) identified above as item(s): 1				
<i>* Where the earlier application is an ARIPO application, it is mandatory to indicate in the Supplemental Box at least one country party to the Paris Convention for the Protection of Industrial Property for which that earlier application was filed (Rule 4.10(b)(iii)). See Supplemental Box.</i>				
Box No. VII INTERNATIONAL SEARCHING AUTHORITY				
Choice of International Searching Authority (ISA) (if two or more International Searching Authorities are competent to carry out the international search, indicate the Authority chosen; the two-letter code may be used):		Request to use results of earlier search; reference to that search (if an earlier search has been carried out by or requested from the International Searching Authority):		
ISA / US		Date (day/month/year)	Number	Country (or regional Office)
Box No. VIII CHECK LIST; LANGUAGE OF FILING				
This international application contains the following number of sheets: request : 4 description (excluding sequence listing part) : 44 claims : 5 abstract : 1 drawings : 1 sequence listing part of description : Total number of sheets : 55		This international application is accompanied by the item(s) marked below: 1. <input checked="" type="checkbox"/> fee calculation sheet 2. <input checked="" type="checkbox"/> separate signed power of attorney 3. <input type="checkbox"/> copy of general power of attorney; reference number, if any: 4. <input type="checkbox"/> statement explaining lack of signature 5. <input type="checkbox"/> priority document(s) identified in Box No. VI as item(s): 6. <input type="checkbox"/> translation of international application into (language): 7. <input type="checkbox"/> separate indications concerning deposited microorganism or other biological material 8. <input type="checkbox"/> nucleotide and/or amino acid sequence listing in computer readable form 9. <input type="checkbox"/> other (specify):		
Figure of the drawings which should accompany the abstract:		Language of filing of the international application: English		
Box No. IX SIGNATURE OF APPLICANT OR AGENT				
Next to each signature, indicate the name of the person signing and the capacity in which the person signs (if such capacity is not obvious from reading the request).				
 WALKER, NAOMI				

For receiving Office use only	
1. Date of actual receipt of the purported international application: 3. Corrected date of actual receipt due to later but timely received papers or drawings completing the purported international application: 4. Date of timely receipt of the required corrections under PCT Article 11(2): 5. International Searching Authority (if two or more are competent): ISA /	2. Drawings: <input type="checkbox"/> received: <input type="checkbox"/> not received: 6. <input type="checkbox"/> Transmittal of search copy delayed until search fee is paid.

For International Bureau use only
Date of receipt of the record copy by the International Bureau:

PCT

FEE CALCULATION SHEET
Annex to the Request

For receiving Office use only

International application No.

Applicant's or agent's
file reference DPL/009/PCT

Date stamp of the receiving Office

Applicant

D-PHARM LIMITED

CALCULATION OF PRESCRIBED FEES

1. TRANSMITTAL FEE ILS 437 T

2. SEARCH FEE USD 700 S

International search to be carried out by ISA/US

(If two or more International Searching Authorities are competent in relation to the international application, indicate the name of the Authority which is chosen to carry out the international search.)

3. INTERNATIONAL FEE

Basic Fee

The international application contains 55 sheets.

first 30 sheets USD 427 b1

25 x 10 - USD 250 b2

remaining sheets additional amount

Add amounts entered at b1 and b2 and enter total at B USD 677 B

Designation Fees

The international application contains _____ designations.

8 x 92 = USD 736 Dnumber of designation fees amount of designation fee
payable (maximum 8)

Add amounts entered at B and D and enter total at I USD 1413 I

(Applicants from certain States are entitled to a reduction of 75% of the international fee. Where the applicant is (or all applicants are) so entitled, the total to be entered at I is 25% of the sum of the amounts entered at B and D.)

4. FEE FOR PRIORITY DOCUMENT (if applicable) ILS 166 P

5. TOTAL FEES PAYABLE ILS 603 USD 2113

Add amounts entered at T, S, I and P, and enter total in the TOTAL box

TOTAL

☐ The designation fees are not paid at this time.

MODE OF PAYMENT

☐ authorization to charge
deposit account (see below)☒ bank draft☐ coupons☐ cheque☐ cash☐ other (specify):☐ postal money order☐ revenue stamps

DEPOSIT ACCOUNT AUTHORIZATION (this mode of payment may not be available at all receiving Offices)

The RO/ ☐ is hereby authorized to charge the total fees indicated above to my deposit account.☐ (this check-box may be marked only if the conditions for deposit accounts of the receiving Office so permit) is hereby authorized to charge any deficiency or credit any overpayment in the total fees indicated above to my deposit account.☐ is hereby authorized to charge the fee for preparation and transmittal of the priority document to the International Bureau of WIPO to my deposit account.

Deposit Account No.

Date (day/month/year)

Signature

PCT

INTERNATIONAL SEARCH REPORT

(PCT Article 18 and Rules 43 and 44)

Applicant's or agent's file reference DPL/009/PCT	FOR FURTHER ACTION	see Notification of Transmittal of International Search Report (Form PCT/ISA/220) as well as, where applicable, item 5 below.
International application No. PCT/IL00/00562	International filing date (<i>day/month/year</i>) 13 September 2000 (13.09.2000)	(Earliest) Priority Date (<i>day/month/year</i>) 14 September 1999 (14.09.1999)
Applicant D-PHARM LIMITED		

This international search report has been prepared by this International Searching Authority and is transmitted to the applicant according to Article 18. A copy is being transmitted to the International Bureau.

This international search report consists of a total of 4 sheets.



It is also accompanied by a copy of each prior art document cited in this report.

1. Basis of the Report

- a. With regard to the **language**, the international search was carried out on the basis of the international application in the language in which it was filed, unless otherwise indicated under this item.



the international search was carried out on the basis of a translation of the international application furnished to this Authority (Rule 23.1(b)).

- b. With regard to any **nucleotide and/or amino acid sequence** disclosed in the international application, the international search was carried out on the basis of the sequence listing:



contained in the international application in written form.



filed together with the international application in computer readable form.



furnished subsequently to this Authority in written form.



furnished subsequently to this Authority in computer readable form.



the statement that the subsequently furnished written sequence listing does not go beyond the disclosure in the international application as filed has been furnished.



the statement that the information recorded in computer readable form is identical to the written sequence listing has been furnished.

2. ☒ **Certain claims were found unsearchable** (See Box I).

3. ☐ **Unity of invention is lacking** (See Box II).

4. With regard to the **title**,



the text is approved as submitted by the applicant.



the text has been established by this Authority to read as follows:

5. With regard to the **abstract**,



the text is approved as submitted by the applicant.



the text has been established, according to Rule 38.2(b), by this Authority as it appears in Box III. The applicant may, within one month from the date of mailing of this international search report, submit comments to this Authority.

6. The figure of the drawings to be published with the abstract is Figure No. 1



as suggested by the applicant.



because the applicant failed to suggest a figure.



because this figure better characterizes the invention.



None of the figures

INTERNATIONAL SEARCH REPORT

Intern application No.

PCT/IL00/00562

Box I Observations where certain claims were found unsearchable (Continuation of Item 1 of first sheet)

This international report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☐ Claim Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:
2. ☐ Claim Nos.:
because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:
3. ☒ Claim Nos.: 14-16
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule

Box II Observations where unity of invention is lacking (Continuation of Item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

☐
☐

The additional search fees were accompanied by the applicant's protest.

No protest accompanied the payment of additional search fees.

Box III TEXT OF THE ABSTRACT (Continuation of Item 5 of the first sheet)**NEW ABSTRACT**

The invention discloses prodrugs comprising anti-proliferative drugs covalently linked, via a bridging group, to a phospholipid moiety such that the active species is preferentially released, preferably by enzymatic cleavage, at the required site of action. The invention further discloses pharmaceutical compositions said prodrugs and the uses thereof for the treatment of diseases and disorders related to inflammatory, to degenerative or atrophic conditions, and to uncontrolled cell growth.

Fig. 1 depicts a graph of animal survival during the course of an experiment wherein mice were i.p. transplanted with 11210 mouse leukemia cells and then treated with vehicle only (squares), MTX (triangles) or molar equivalent dose of DP-MTX071 (circles) according to the regimen described in example in Example 11.

INTERNATIONAL SEARCH REPORT

International application No.

PCT/IL00/00562

A. CLASSIFICATION OF SUBJECT MATTER

IPC(7) : A61K 31/56

US CL : 514/178, 180, 169, and 424/450

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

U.S. : 514/178, 180, 169, and 424/450

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y ---- A	US 4,772, 594 A (HASHIMOTO et al) 20 September 1988, see col. 1, lines 13-48 cols.3, line 67-col. 6, line 14.	1,3,17 ----- 2, 4-13, and 18-29
Y,P	US 6,028,066 A (UNGER) 22 February 2000, see entire document especially col.1,line 54to col.3, line 14, col.11, line 23-52, col. 18, line 45-60, col.23, line 36 -col 25, line 20 example 10.	1-13, 17-29
Y,P	US 6,077,837 A (KOZAK) 20 June 2000, see entire document.	1-13, 17-29

☐ Further documents are listed in the continuation of Box C.☐ See patent family annex.

* Special categories of cited documents:

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier application or patent published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T"

later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X"

document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y"

document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art

"&"

document member of the same patent family

Date of the actual completion of the international search

19 April 2001 (19.04.2001)

Date of mailing of the international search report

30 APR 2001

Name and mailing address of the ISA/US

Commissioner of Patents and Trademarks
Box PCT
Washington, D.C. 20231

Facsimile No. (703)305-3230

Authorized officer

SHEP ROSE

Telephone No. (703) 308-0193

INTERNATIONAL SEARCH REPORT

International application No.

PCT/IL00/00562

A. CLASSIFICATION OF SUBJECT MATTER

IPC(7) : A61K 31/56

US CL : 514/178, 180, 169, and 424/450

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

U.S. : 514/178, 180, 169, and 424/450

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y — A	US 4,772, 594 A (HASHIMOTO et al) 20 September 1988, see col. 1, lines 13-48 cols.3, line 67-col. 6, line 14.	1,3,17 2, 4-13, and 18-29
Y,P	US 6,028,066 A (UNGER) 22 February 2000, see entire document especially col.1, line 54 to col.3, line 14, col.11, line 23-52, col. 18, line 45-60, col.23, line 36 -col 25, line 20 example 10.	1-13, 17-29
Y,P	US 6,077,837 A (KOZAK) 20 June 2000, see entire document.	1-13, 17-29

☐ Further documents are listed in the continuation of Box C.

☐ See patent family annex.

* Special categories of cited documents:

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier application or patent published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T"

later documents published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X"

document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y"

document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art

"&"

document member of the same patent family

Date of the actual completion of the international search

19 April 2001 (19.04.2001)

Date of mailing of the international search report

30 APR 2001

Name and mailing address of the ISA/US

Commissioner of Patents and Trademarks

Box PCT

Washington, D.C. 20231

Facsimile No. (703)305-3230

Authorized officer

SHEP ROSE

Telephone No. (703) 308-0193

INTERNATIONAL SEARCH REPORT

Inter. onal application No.

PCT/IL00/00562

Box I Observations where certain claims were found unsearchable (Continuation of Item 1 of first sheet)

This international report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☐ Claim Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:
2. ☐ Claim Nos.:
because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:
3. ☒ Claim Nos.: 14-16
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule

Box II Observations where unity of invention is lacking (Continuation of Item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

☐
☐

The additional search fees were accompanied by the applicant's protest.

No protest accompanied the payment of additional search fees.

From the
INTERNATIONAL PRELIMINARY EXAMINING AUTHORITY

To:
NAOMI WALKER
KIRYAT WEIZMANN SCIENCE PARK
P.O. BOX 2313
REHOVOT, ISRAEL 76123

PCT

NOTIFICATION OF RECEIPT
OF DEMAND BY COMPETENT INTERNATIONAL
PRELIMINARY EXAMINING AUTHORITY

(PCT Rules 59.3(e) and 61.1(b), first sentence
and Administrative Instructions, Section 601(a))

Date of mailing
(day/month/year)

10 JUL 2001

Applicant's or agent's file reference

DPL/009/PCT

IMPORTANT NOTIFICATION

International application No.

PCT/IL00/00562

International filing date (day/month/year)

13 SEP 00

Priority date (day/month/year)

14 SEP 99

Applicant

D-PHARM LIMITED

1. The applicant is hereby notified that this International Preliminary Examining Authority considers the following date as the date of receipt of the demand for international preliminary examination of the international application:

06 APR 2001

2. That date of receipt is:



the actual date of receipt of the demand by this Authority (Rule 61.1(b)).



the actual date of receipt of the demand on behalf of this Authority (Rule 59.3(e)).



the date on which this Authority has, in response to the invitation to correct defects in the demand (Form PCT/IPEA/404), received the required corrections.

3. ☐ **ATTENTION:** That date of receipt is **AFTER** the expiration of 19 months from the priority date. Consequently, the election(s) made in the demand does (do) not have the effect of postponing the entry into the national phase until 30 months from the priority date (or later in some Offices) (Article 39(1)). Therefore, the acts for entry into the national phase must be performed within 20 months from the priority date (or later in some Offices) (Article 22). For details, see the *PCT Applicant's Guide, Volume II*.



(If applicable) This notification confirms the information given by telephone, facsimile transmission or in person on:

4. Only where paragraph 3 applies, a copy of this notification has been sent to the International Bureau.

Name and mailing address of the IPEA/
Assistant Commissioner for Patent
Box PCT
Washington, D.C. 20231 Attn:RO/US
Facsimile No. 703-305-3230

Authorized officer

Rayna Lader
Telephone No. (703) 305-5297